

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : G01N 31/00	A1	(11) International Publication Number: WO 99/12027 (43) International Publication Date: 11 March 1999 (11.03.99)
(21) International Application Number: PCT/US98/12351 (22) International Filing Date: 12 June 1998 (12.06.98) (30) Priority Data: 60/057,411 29 August 1997 (29.08.97) US (71) Applicant (for all designated States except US): THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; Fifth floor, 1111 Franklin Street, Oakland, CA 94607-5200 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): REICH, Norbert, O. [US/US]; 450 San Domingo Drive, Santa Barbara, CA 93111 (US). FLYNN, James [US/US]; 5658 Armitos Avenue, Goleta, CA 93117 (US). (74) Agent: CANADY, Karen, S.; Merchant, Gould, Smith, Edell, Welter & Schmidt, Suite 400, 11150 Santa Monica Boulevard, Los Angeles, CA 90025-3395 (US).		(81) Designated States: CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: MODULATORS OF DNA CYTOSINE-5 METHYLTRANSFERASE AND METHODS FOR USE THEREOF		
(57) Abstract A synthetic oligonucleotide comprising a C-5 methylcytosine and which recognizes and binds an allosteric site on DNA methyltransferase thereby inhibiting DNA methyltransferase activity is disclosed. Also disclosed is a composition comprising a synthetic oligonucleotide of the invention. The composition is useful for inhibiting DNA methyltransferase activity, thereby inhibiting the methylation of DNA. The composition can be a pharmaceutical composition useful for treating disorders associated with methylation defects, such as cancer and certain developmental disorders. Also disclosed is a method of inhibiting methylation of DNA. The method involves contacting a DCMTase with a synthetic oligonucleotide of the invention in the presence of the DNA, thereby resulting in an enzyme/synthetic oligonucleotide complex. The presence of the complex prevents catalysis, thereby inhibiting DNA methyltransferase activity. Also disclosed is a method of treating a disorder of cell proliferation or development by administering to a subject a synthetic oligonucleotide of the invention. The inhibition of DNA methyltransferase prevents the methylation of DNA thereby treating the disorder of cell proliferation or development.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

MODULATORS OF DNA CYTOSINE-5 METHYLTRANSFERASE
AND METHODS FOR USE THEREOF

This application is based on United States provisional patent application serial number 60/057,411, filed August 29, 1997, the entire contents of which are hereby incorporated by reference into this application. Throughout this application various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

- 10 This invention was made with Government support under Grant No. GM46333, awarded by the National Institutes of Health to Norbert O. Reich. The Government has certain rights in this invention.

BACKGROUND OF THE INVENTION

In eukaryotic organisms, DNA methylation is catalyzed by an S-adenosyl-L-methionine (AdoMet)¹-dependent DNA cytosine-C⁵ methyltransferase (DCMTase, EC 2.1.1.37). Methyl group transfer to the cytosine-C⁵ position occurs predominately within the cytosyl-guanosyl (CpG) context (Boyes, J., & Bird, A.P., 1991, DNA methylation inhibits transcription indirectly via a methyl-CpG binding protein, Cell 64:1123-1134). The genomic distribution of 5-methylcytosine (5-mC) dynamically changes throughout ontogeny (Razin, A., & Riggs, A.D., 1980, DNA methylation and gene function, Science 210:604-609; Kafri, T. et al., 1992, Developmental pattern of gene-specific DNA methylation in the mouse embryo and germ line, Genes and Dev. 6:705-714). The methylation state of a gene specifically affects transcription.

- 25 DCMTase is involved in mammalian development by way of an undefined process that can lead to gene regulation (reviewed in Jost, J.P., & Saluz, H.P., 1993, DNA Methylation: Molecular Biology and Biological Significance, Birkhauser Verlag, Basel). Proper DCMTase function is essential for viable development and for normal cellular activity (Li, E. et al., 1992, Targeted mutation of the DNA methyltransferase gene results in embryonic lethality, Cell 69:915-926).

Cytosine methylation is the predominant epigenetic event in the modification of eukaryotic DNA. To date only a single DCMTase has been identified in several metazoan organisms (Yoder, J.A., et al., 1996, New 5' regions of the murine and

human genes for DNA cytosine-5 methyltransferase, J. Biol. Chem. 271:31092-31097). The function most often identified with cytosine C⁵ methylation (5-^mC) in higher eukaryotes is the regulation of transcription (Jost, J.P., & Saluz, H.P., 1993, DNA Methylation: Molecular Biology and Biological Significance, Birkhauser Verlag, Basel). Generally, hypermethylated genes are transcriptionally silent and inheritance of the proper genomic methylation pattern is critical to viable development as shown by DCMTase gene knock-outs in mice (Li, E., et al., 1992, Targeted mutation of the DNA methyltransferase gene results in embryonic lethality, Cell 69:15-926). Anti-sense directed inactivation of DCMTase mRNA as well as the incorporation of the cytosine analogs 5-azacytidine and 5-fluorocytidine into DNA interfere with DCMTase function and lead to cytological dysfunction (Ramachandani, S., et al., 1997, Inhibition of tumorigenesis by a cytosine-DNA, methyltransferase, antisense oligodeoxynucleotide, Proc. Natl. Acad. Sci. USA 94:684-689; Jones, P.A., 1985, Altering gene expression with 5-azacytidine, Cell 40:485-486).

Eukaryotic DCMTase cDNAs have been cloned and sequenced; five are from animal sources (mouse: Bestor, T., et al., 1988, Cloning and sequencing of a cDNA encoding DNA methyltransferase of mouse cells, J. Mol. Biol. 203:971-983; human: Yen, R.C., et al., 1992, Isolation and characterization of the cDNA encoding human DNA methyltransferase, Nucleic Acids Res. 20:2287-2291; chicken: Tajima, S., et al., 1995, Isolation and expression of a chicken DNA methyltransferase cDNA, J. Biochem. 117:1050-1057; frog: Kimura et al., 1996, Isolation and expression of a *Xenopus laevis* DNA methyltransferase cDNA, Journal of Biochemistry, 120:1182-1189; sea urchin: Aniello et al., 1996, Isolation of cDNA clones encoding DNA methyltransferase of sea urchin *P. lividus*: expression during embryonic development, Gene 178:57-61). These DCMTases are composed of a large amino-terminal domain and a smaller carboxy-terminal domain that contains many of the major motifs found in prokaryotic DCMTases (Posfai, J., et al., 1989, Predictive motifs derived from cytosine methyltransferases, Nucleic Acids Res. 17:2421-2435). The amino-terminal domain has been implicated in nuclear localization to DNA replication foci during S-phase (Leonhardt, H., et al., 1992, A targeting sequence directs DNA methyltransferase to sites of DNA replication in mammalian nuclei, Cell 71:865-873), metal binding by zinc finger domains, and DNA binding (Bestor, T.H., 1992, Activation of the mammalian DNA methyltransferase by cleavage of a Zn binding regulatory domain, EMBO 11:2611-2617; Chuang, L.S., et al., 1996, Characterisation of independent DNA and multiple Zn-binding domains at the N terminus of human

DNA-(cytosine-5) methyltransferase: modulating the property of a DNA-binding domain by contiguous Zn-binding motifs, Chia, J., and Li, B.F.L., J. Mol. Biol. 257:935-948).

- 5 Although the cellular processes that determine the genomic patterns of DNA methylation are not understood, DCMTase has an essential role in these processes. A basic understanding of the binding and catalytic DNA sequence specificity (discrimination) of the enzyme, and the factors which regulate this specificity are important. Since the mammalian enzyme is a relatively large, 183 kDa protein, DNA
- 10 sequences flanking the cognate CpG may modulate the ability of the enzyme to methylate particular CpG sites. However, the CpG flanking sequence preferences of the enzyme, and its preference for single- and double-stranded substrates have not been rigorously addressed by previous investigators (Bestor, T.H. et al., 1992, CpG islands in mammalian gene promoters are inherently resistant to de novo methylation,
- 15 GATA 9:48-53; Hepburn, P.A., et al., 1991, Enzymatic methylation of cytosine in DNA is prevented by adjacent O⁶-methylguanine residues, J. Biol. Chem. 266:7985-7987; Bolden, A.H., et al., 1986, Primary DNA sequence determines sites of maintenance and de novo methylation by mammalian DNA methyltransferases, Mol. Cell. Bio. 6:1135-1140; Pfeifer, G.P., et al., 1985, Mouse DNA-cytosine-5-
- 20 methyltransferase: sequence specificity of the methylation reaction and electron microscopy of enzyme-DNA complexes, EMBO J. 4:2879-2884; Ward, C., et al., 1987, *In vitro* methylation of the 5'-flanking regions of the mouse b-globin gene, J. Biol. Chem. 262:11057-11063; Carotti, D., et al., 1986, Substrate preferences of the human placental DNA methyltransferase investigated with synthetic
- 25 polydeoxynucleotides, Biochim. et Biophys. Acta. 866:135-143; Carotti D. et al., 1986, *supra*; Wang, R.Y.H., et al., 1984, Human placental DNA methyltransferase: DNA substrate and DNA binding specificity, Nucl. Acids Res. 12:3473-3490; Pfeifer et al., 1985, *supra*; Gruenbaum, Y., et al., 1982, Substrate and sequence specificity of a eukaryotic DNA methylase, Nature 295:620-622).

30

- There is evidence that errors in the proper maintenance of genomic methylation are involved in aging and cancer. CpG islands are reported to become hypermethylated with age and may down-regulate expression of essential genes (Antequerra & Bird, 1993, Number of CpG islands and genes in human and mouse, Proceedings of the
- 35 National Academy of Sciences, USA, 90:11995-11999; Nyce, J.W., 1997, Drug-induced DNA hypermethylation: A potential mediator of acquired drug resistance

during cancer chemotherapy, Mutation Research 386:153-161) Amplification of DCMTase expression by an exogenous mammalian DCMTase gene induces tumorigenic transformation of NIH 3T3 mouse fibroblasts (Wu et al., 1993, Expression of an exogenous eukaryotic DNA methyltransferase gene induces transformation of NIH 3T3 cells, Proc. Natl. Acad. Sci., USA, 90:8891-8895). Human neoplastic cells and cells derived from different stages of colon cancer express up to 200-fold higher levels of DCMTase than normal (El-Deiry et al., 1991, High expression of the DNA methyltransferase gene characterizes human neoplastic cells and progression stages of colon cancer, Proc. Natl. Acad. Sci., USA, 88:3470-3474). This contributes substantially to tumor development in a mouse model of intestinal neoplasia (Laird, P.W., et al., 1995, Suppression of intestinal neoplasia by DNA hypomethylation, Cell 81:197-205). Changes in DNA methylation and DCMTase activity appear early in oncogenesis (Belinsky, S.A., et al., 1996, Increased cytosine DNA-methyltransferase activity is target-cell-specific and an early event in lung cancer, Proc. Natl. Acad. Sci. USA 93:4045-4050).

Conversely, antisense oligonucleotides that interfere with expression of DCMTase may inhibit tumorigenesis (Ramachandani et al., 1997, Inhibition of tumorigenesis by a cytosine-DNA methyltransferase, antisense oligonucleotide, Proc. Natl. Acad. Sci., USA, 94:684-689; MacLeod & Szyf, 1995, Expression of antisense to DNA methyltransferase mRNA induces DNA demethylation and inhibits tumorigenesis, J. Biol. Chem. 270:8037-8043). The anticancer agent 5-aza-deoxycytidine functions by inhibiting the DCMTase (Jones, 1985, Altering gene expression with 5-azacytidine, Cell 40:485-486; Jutterman et al., 1994, Toxicity of 5-aza-2'-deoxycytidine to mammalian cells is mediated primarily by covalent trapping of DNA methyltransferase rather than DNA demethylation, Proc. Natl. Acad. Sci., USA, 91:11797-11801). Changes in DNA methylation and DCMTase activity early in oncogenesis (Belinsky, S.A., et al., 1996, *supra*) and the ability of DCMTase inhibitors to virtually abolish adenoma formation in mice (Laird, P.W., et al., 1995, *supra*) suggest that DCMTase inhibitors might be useful anticancer therapeutics (Szyf, M., 1996, The DNA methylation machinery as a target for anticancer therapy, Pharmacol. Ther. 70:1-37). 5-Aza-deoxycytidine is an irreversible, mechanism-based DCMTase inhibitor that has been used in patients with acute myeloid leukemia. Unfortunately, 5-Aza-deoxycytidine is unstable in solution and may be carcinogenic as well as mutagenic (Jones, P.A., 1996, DNA methylation errors and cancer, Cancer Res. 56:2463-2467). There is a need for DCMTase inhibitors that do not require

incorporation into DNA and that are mechanistically unlike 5-aza-deoxycytidine (Belinsky, S.A., et al., 1996, *supra*; Szyf, M., 1996, *supra*; Jones, 1996, *supra*). A keen understanding of how DCMTase functions *in vitro* can be the basis for better strategies to both activate and inhibit the enzyme to correct developmental disorders like cancer.

Enzymes that catalyze one carbon additions to C⁵ of pyrimidines define a class of enzymes with similar chemistry (Ivanetich, K.M., & Santi, D.V., 1992, 5,6-Dihydropyrimidine adducts in the reactions and interactions of pyrimidines with proteins, *Prog. Nucleic Acid Res. Mol. Biol.* 42:127-156). The bacterial DNA cytosine C⁵ methyltransferase, *M.HhaI* (38 kDa Mr), modifies the internal cytosine in GCGC and has an ordered Bi Bi kinetic mechanism in which DNA binds first (Wu, J.C., & Santi, D.V., 1987, Kinetic and catalytic mechanism of *HhaI* methyltransferase, *J. Biol. Chem.* 262:4778-4786). Catalysis involves nucleophilic attack of an active site cysteine at the C⁶ position of the cytosine which, in the absence of the cofactor, leads to exchange of the C⁵ hydrogen. A *M.HhaI*-DNA cocrystal structure suggests that a catalytic intermediate exists that involves the translocation of the target cytosine to an extrahelical position covalently bound to an active site cysteine (Klimasauskas, S., et al., 1994. *HhaI* methyltransferase flips its target base out of the DNA helix, *Cell* 76:357-369). Methyl transfer from AdoMet is followed by β -elimination to regenerate the active enzyme (Wu & Santi, 1987, *supra*; Osterman, D.G., et al., 1988, 5-Fluorocytosine in DNA is a mechanism-based inhibitor of *HhaI* methylase, *Biochemistry* 27:5204-5210).

A recent kinetic study of a highly homogeneous, unproteolyzed preparation of DCMTase from mouse erythroleukemia cells (MEL) further characterized the interactions of the enzyme with DNA and AdoMet (Flynn, J., et al., 1996, Murine DNA cytosine-C5 methyltransferase: Pre-steady- and steady-state kinetic analyses with regulatory DNA sequences, *Biochemistry* 35:7308-7315). The invention disclosed herein descriptively accounts for the previously reported complexities in kinetic behavior and identifies a potent single-stranded oligonucleotide inhibitor that binds to the enzyme at a distinct regulatory site.

There is a need for molecules which modulate the methylation of DNA for the reasons discussed above. In addition, molecules which inhibit DNA methylation can be useful for preventing drug resistance acquired by subjects undergoing cancer chemotherapy.

Drug-induced DNA hypermethylation is regarded as a potential mediator of this acquired drug resistance (Nyce, J.W., 1997, Drug-induced DNA hypermethylation: A potential mediator of acquired drug resistance during cancer chemotherapy, Mutation Research 386:153-161).

5

SUMMARY OF THE INVENTION

The invention provides synthetic oligonucleotides comprising a C-5 methylcytosine. The oligonucleotide recognizes and binds an allosteric site on DNA methyltransferase thereby inhibiting DNA methyltransferase activity. In one embodiment, the synthetic
10 oligonucleotide has an inhibition constant of not greater than 1000 nM. In another embodiment, the synthetic oligonucleotide has an inhibition constant of not greater than 200 nM. In yet another embodiment, the synthetic oligonucleotide has an inhibition constant of not greater than 20 nM.

15 The invention further provides a composition comprising a synthetic oligonucleotide comprising a C-5 methylcytosine and which recognizes and binds an allosteric site on DNA methyltransferase. The composition is useful for inhibiting DNA methyltransferase activity, thereby inhibiting the methylation of DNA. In one embodiment, the composition is a pharmaceutical composition comprising a
20 pharmaceutically effective amount of a synthetic oligonucleotide comprising a C-5 methylcytosine and which recognizes and binds an allosteric site on DNA methyltransferase, and optionally, a pharmaceutically acceptable carrier. The pharmaceutical composition is useful for treating disorders associated with methylation defects, such as cancer and certain developmental disorders.

25

The invention further provides a method of inhibiting methylation of DNA. The method involves contacting a DCMTase with a synthetic oligonucleotide which recognizes and binds an allosteric site on DNA methyltransferase thereby resulting in a DNA methyltransferase/synthetic oligonucleotide complex. The complex is contacted with the
30 DNA. The presence of the complex prevents binding of AdoMet to DNA methyltransferase in a catalytically competent manner thereby inhibiting DNA methyltransferase activity and inhibiting methylation of DNA. In one embodiment, the synthetic oligonucleotide comprises a C-5 methylcytosine.

35 The invention further provides a method of treating a disorder of cell proliferation or development. The method involves administering to a subject a synthetic

- oligonucleotide which recognizes and binds an allosteric site on DNA methyltransferase. The binding of the synthetic oligonucleotide prevents binding of AdoMet to DNA methyltransferase in a catalytically competent manner thereby inhibiting DNA methyltransferase. The inhibition of DNA methyltransferase prevents the methylation of DNA thereby treating the disorder of cell proliferation or development. In one embodiment, the synthetic oligonucleotide comprises a C-5 methylcytosine. In one embodiment, the disorder of cell proliferation is cancer such as lung cancer, breast cancer, prostate cancer, pancreatic cancer or colon cancer.
- 10 The invention also provides a method of identifying a modulator of DCMTase which recognizes and binds an allosteric site on DCMTase. The method comprises contacting a molecule with DCMTase in the presence of AdoMet and DNA. The method further comprises measuring DCMTase activity. An increase or decrease in DCMTase activity is indicative of a modulator of DCMTase activity. In one embodiment, the modulator is an inhibitor. In another embodiment, the modulator is an activator.

BRIEF DESCRIPTION OF THE FIGURES

- Figure 1A shows six synthetic oligonucleotides that mimic the GC-box and the cyclic AMP responsive elements (CRE) (SEQ ID NOS:9-12). The appropriate consensus is in bold type and the single, centrally located CpG dinucleotide is underlined (mC = C-5 methylcytosine). The complementary a, aMET, b, and bMET strands were annealed to produce unmethylated, a/b, and hemi-methylated, aMET/b or a/bMET, double-stranded substrates.

- 25 Figure 1B shows oligonucleotide sequences corresponding to SEQ ID NOS:10, 11, 13, 14 and 15, as indicated, which were tested for inhibition in an in vitro assay. GC-box pMET has a phosphorothioate backbone, while the others have a deoxyribose backbone. K_{ij} is the inhibition constant derived from the y-intercept values from double reciprocal plots and is a characteristic of the inhibitor binding to the allosteric site of the enzyme. IC_{50} is the concentration of inhibitor that produces 50% activity of an uninhibited reaction.

- Figure 2 is an autoradiogram showing the results of gel mobility shift analysis varying DCMTase with constant GC-box a/b. Lane 1: 0 nM DCMTase; Lane 2: 5.0 nM DCMTase; Lane 3: 10 nM DCMTase; Lane 4: 20 nM DCMTase; Lane 5: 30 nM DCMTase; Lane 6: 35 nM DCMTase; Lane 7: 40 nM DCMTase; Lane 8: 45 nM

DCMTase; Lane 9: 50 nM DCMTase; Lane 10: 65 nM DCMTase; Lane 11: 75 nM DCMTase; Lane 12: 95 nM DCMTase.

Figure 3 is an autoradiogram showing the results of a gel mobility shift analysis
 5 varying GC-box a/b^{MET} with constant DCMTase. Lane 1: 0.050 μ M Free DNA; Lane 2: 0.10 μ M Free DNA; Lane 3: 0.15 μ M Free DNA; Lane 4: 0.10 μ M DNA; Lane 5: 0.28 μ M DNA; Lane 6: 0.45 μ M DNA; Lane 7: 0.63 μ M DNA; Lane 8: 0.80 μ M DNA; Lane 9: 1.0 μ M DNA; Lane 10: 2.0 μ M DNA; Lane 11: 2.0 μ M Free DNA. Lanes 1, 2, 3, and 11 are control experiments without added DCMTase.

10 Figure 4 is an autoradiogram showing the results of a gel mobility shift analysis varying GC-box a/b with constant DCMTase. Lane 1: 0.050 μ M Free DNA; Lane 2: 0.10 μ M Free DNA; Lane 3: 0.15 μ M Free DNA; Lane 4: 0.10 μ M DNA; Lane 5: 0.50 μ M DNA; Lane 6: 1.0 μ M DNA; Lane 7: 4.0 μ M DNA; Lane 8: 6.0 μ M DNA;
 15 Lane 9: 6.0 μ M Free DNA. Lanes 1, 2, 3, and 9 are control experiments without added DCMTase.

Figure 5 is an autoradiogram showing the results of a gel mobility shift analysis
 20 varying GC-box b^{MET} with constant DCMTase. Lane 1: 0.10 μ M Free DNA; Lane 2: 0.20 μ M DNA; Lane 3: 0.40 μ M DNA; Lane 4: 0.80 μ M DNA; Lane 5: 1.6 μ M DNA; Lane 6: 3.2 μ M DNA; Lane 7: 6.4 μ M DNA; Lane 8: 3.2 μ M DNA; Lane 9: 6.0 μ M CRE a^{MET}/b. Lanes 1 and 8 are control experiments without added DCMTase.

Figure 6 shows a randomized DNA substrate used in *in vitro* screening (SEQ ID
 25 NOS:16-17). The top strand shown was synthesized using b-cyanoethyl phosphoramidite chemistry. The PCR primers used for amplifying the shifted DNA are underlined. Primer C is underlined and contains an EcoRI restriction site. Primer D, underlined twice, contains a BamHI restriction site and was annealed to the randomized top strand for extension by Klenow polymerase. The randomized positions
 30 are denoted as N and are either dG, dA or dT on one strand and the complementary dC, dA or dT on the other strand of the duplex.

Figure 7 shows cloned and sequenced individual isolates from the pooled generations (SEQ ID NOS:18-100 respectively). Only the guanine containing strand is shown for
 35 simplicity. Generation-5 members are arranged with the highest guanine content on

the 5' side of the invariant CpG at the top. Frequency information is given for each randomized flank on the appropriate border, an asterisk denotes a single occurrence.

5 Figure 8A shows the nucleotide frequency at each randomized flanking position for the generation-5 screening in the form of a bar graph indicating the percent occurrence of each nucleotide at the randomized positions. The predominance of guanosine extends over the entire randomized region. The horizontal line at 33% is representative of the starting pool frequencies. The line at 70% is added as a visual aid.

10

Figure 8B lists the nucleotide percentages at each randomized position for the generation-5 screening.

15 Figure 9 shows genomic sequences similar to the DCMTase selected generation-5 clones (SEQ ID NOS:101-110). Fasta searches through the mouse and human GenBank libraries produced these matches when limited to no greater than four mismatches and no gaps. The definitions have been edited from the original entries.

20 Figure 10 shows initial velocity curves of the selected generations. Squares, generation-1 pool; triangles, generation-2 pool; circles, generation-4 pool; diamonds, generation-5 pool.

25 Figure 11 shows substrate inhibition plots. Reactions contained 3.0 nM DCMTase and 10 μ M AdoMet in MR buffer. The inset shows data in which GC-box a/b was the substrate, using 100 nM DCMTase. Experimental data are shown scattered around a line fit to equation 1 for substrate inhibition. For a direct comparison of the DNA substrates, data are expressed as a V_{max} normalized, S/K_m^{DNA} ratio.

30 Figure 12A shows double reciprocal plots of velocity versus substrate concentration. Poly(dI-dC:dI-dC) was varied and lines represent a constant AdoMet concentration: triangles, 4 μ M; squares, 2 μ M; diamonds, 1 μ M; circles, 0.5 μ M. Experimental data are shown scattered around lines derived from the fit of equation 2 for a sequential mechanism.

35 Figure 12B shows double reciprocal plots of velocity versus substrate concentration. AdoMet was varied and lines represent a constant poly(dI-dC:dI-dC) concentration:

triangles, 112 pM; squares, 56 pM; diamonds, 28 pM; circles, 14 pM. Experimental data are shown scattered around lines derived from the fit of equation 2 for a sequential mechanism.

- 5 Figure 13 shows a double reciprocal plot of velocity versus poly(dI·dC:dI·dC) with varying GC-box b concentrations. The GC-box b concentrations were: diamonds, 0; circles, 0.75 μ M; triangles, 1.5 μ M; squares, 5.0 μ M. Experimental data are shown scattered around lines derived from the fit to equation 5 for noncompetitive inhibition.
- 10 Figure 14 is a double reciprocal plot of velocity vs. poly(dI·dC:dI·dC) with varying GC-box b^{MET} concentrations. The GC-box b^{MET} concentrations were: squares, 0; circles, 10 nM; diamonds, 20 nM; triangles, 40 nM. Experimental data are shown scattered around lines derived from a fit to the log form of equation 6 for uncompetitive inhibition.
- 15 Figure 15 shows a double reciprocal plot of velocity versus AdoMet with varying GC-box b^{MET} concentrations. The GC-box b^{MET} concentrations were: squares, 0; circles, 20 nM; diamonds, 40 nM; triangles, 80 nM. Experimental data are shown scattered around lines derived from a fit to equation 4 for competitive inhibition.
- 20 Figure 16 shows a double reciprocal plot of AdoHcy product inhibition with varying AdoMet concentrations. The AdoHcy concentrations were: squares, 0; diamonds, 0.75 μ M; circles, 1.5 μ M; triangles, 3.0 μ M; notched squares, 6.0 μ M. Experimental data are shown scattered around lines derived from a fit to equation 4 for competitive
- 25 inhibition.
- Figure 17A shows a double reciprocal plot of AdoHcy product inhibition with varying poly(dI·dC:dI·dC) concentrations, in which AdoMet was held constant at 1.2 μ M. The AdoHcy concentrations were: squares, 0; diamonds, 15 μ M; circles, 30 μ M.
- 30 Figure 17B shows a double reciprocal plot of AdoHcy product inhibition with varying poly(dI·dC:dI·dC) concentrations, in which AdoMet was held constant at 8 μ M. The AdoHcy concentrations were: squares, 0; diamonds, 15 μ M; circles, 30 μ M.
- 35 Figure 17C shows a double reciprocal plot of AdoHcy product inhibition with varying poly(dI·dC:dI·dC) concentrations. The AdoHcy concentrations were: squares, 0;

diamonds, 15 μM ; circles, 30 μM . These are secondary slope replots from a series of experiments in which the AdoMet concentrations were: circles, 6.3 μM ; diamonds, 2.5 μM ; squares 1 μM .

5 Figure 18 shows a double reciprocal plot of poly(dId^mC:dId^mC) product inhibition with varying AdoMet concentrations. The poly(dId^mC:dId^mC) concentrations were: squares, 0; diamonds, 5.0 pM; circles, 10 pM; triangles, 20 pM. Experimental data are shown scattered around lines derived from a fit to equation 5 for noncompetitive inhibition.

10

Figure 19A shows a double reciprocal plot of poly(dId^mC:dId^mC) product inhibition with varying poly(dI·dC:dI·dC) concentrations. The poly(dId^mC:dId^mC) concentrations were: squares, 0; triangles, 34 pM; circles, 45 pM; diamonds, 68, notched squares, 90 pM. Experimental data are shown scattered around lines derived from a fit to equation 4 for competitive inhibition. The fitting is not acceptable.

15

Figure 19B shows a double reciprocal plot of poly(dId^mC:dId^mC) product inhibition with varying poly(dI·dC:dI·dC) concentrations. The poly(dId^mC:dId^mC) concentrations were: squares, 0; triangles, 34 pM; circles, 45 pM; diamonds, 68, notched squares, 90 pM. Experimental data are shown scattered around lines derived from a fit to equation 5 for noncompetitive inhibition. The fitting is not acceptable.

20

Figure 20 shows initial velocity plots of different poly(dIdC:dIdC) lengths. The poly(dI·dC:dI·dC) sizes were: circles, 100 base-pairs; diamonds, 500 base-pairs; triangles, 2000 base-pairs; squares, 5000 base-pairs. The inset provides a zoom in along the x-axis toward the origin to show the quality of the data.

25

Figure 21 shows a plot of DCMTase specificity as a function of poly(dIdC:dIdC) length. The apparent constants were derived from Figure 20 and are shown in Table 6. The data was fit well by an isotherm that yielded a half-maximal length of 1200 base-pairs and a maximal specificity value of $29 \times 10^{11} \text{ hr}^{-1} \text{ pM}^{-1}$ with poly(dIdC:dIdC) as the substrate.

30

Figure 22 shows a proposed kinetic mechanism. DCMTase appears to progress through the catalytic cycle by the Ordered Bi-Bi mechanism shown.

35

Figure 23A is a double-reciprocal plot of poly(dId^mC:dId^mC) product inhibition with varying poly(dI-dC:dI-dC) concentrations. Reactions contained 20 nM DCMTase and 1.5 μ M AdoMet in 100 mM Tris pH 8.0, 10 mM EDTA, 10 mM DTT, 200 μ g/mL BSA. Incubations were at 37 °C for 60 minutes. The poly(dI-dC:dI-dC) concentrations were 20, 40, 80, 120 and 160 pM. The poly(dId^mC:dId^mC) concentrations were: squares, 0; triangles, 34 pM; circles, 45 pM; diamonds, 68, notched squares, 90 pM. Shown are the intersecting noncompetitive lines.

Figure 23B is the slope replot of the plot shown in Figure 23A.

Figure 23C is the y-intercept replot obtained from the lines in Figure 23A.

10 DETAILED DESCRIPTION OF THE INVENTION

The invention provides a synthetic oligonucleotide comprising a C-5 methylcytosine and which recognizes and binds an allosteric site on DNA cytosine methyltransferase thereby modulating DCMTase activity associated with the allosteric site. In one embodiment, the modulating comprises inhibition. In another embodiment, the modulating comprises activation. The C-5 methylcytosine of the synthetic oligonucleotide can be present as a 5mCpG dinucleotide.

In one embodiment, the DCMTase is from a mammal, bird, fish, amphibian, reptile, insect, plant, bacterium, virus or fungus. The mammal can be selected from the group consisting of mouse and human.

In one embodiment, the synthetic oligonucleotide comprises a nucleotide sequence as shown in Figure 1B and designated GC-box b^{MET} (SEQ ID NO:10), GC-box p^{MET} (SEQ ID NO:10), GC-box c^{MET} (SEQ ID NO:13), GC-box d^{MET} (SEQ ID NO:14), GC-box e^{MET} (SEQ ID NO:15), or CRE a^{MET} (SEQ ID NO:11). In one embodiment, the synthetic oligonucleotide has an inhibition constant of not greater than 1000 nM by steady-state kinetic assay. In another embodiment, the synthetic oligonucleotide has an inhibition constant of not greater than 200 nM by steady-state kinetic assay. In yet another embodiment, the synthetic oligonucleotide has an inhibition constant of not greater than 20 nM by steady-state kinetic assay.

In accordance with the practice of the invention, the oligonucleotide can be DNA, RNA, or a derivative or hybrid thereof. The invention further provides a composition comprising a synthetic oligonucleotide comprising a C-5 methylcytosine and which

recognizes and binds an allosteric site on DNA methyltransferase. The composition is useful for inhibiting DNA methyltransferase activity, thereby inhibiting the methylation of DNA. In one embodiment, the composition is a pharmaceutical composition comprising a pharmaceutically effective amount of a synthetic oligonucleotide
5 comprising a C-5 methylcytosine, or a pharmaceutically acceptable salt thereof, and which recognizes and binds an allosteric site on DNA methyltransferase. In one embodiment, the pharmaceutical composition further comprises a pharmaceutically acceptable carrier. The pharmaceutical composition is useful for treating disorders associated with methylation defects, such as cancer and certain developmental disorders.

10

The invention further provides a method of inhibiting methylation of DNA. The method involves contacting a DNA methyltransferase with a synthetic oligonucleotide which recognizes and binds an allosteric site on DNA methyltransferase thereby resulting in an enzyme/synthetic oligonucleotide complex. The presence of the complex prevents
15 binding of AdoMet to DNA methyltransferase in a catalytically competent manner thereby inhibiting DNA methyltransferase activity and inhibiting methylation of DNA. In one embodiment, the enzyme/synthetic oligonucleotide complex forms a further complex with DNA. In one embodiment, the synthetic oligonucleotide comprises a C-5 methylcytosine. In one embodiment, the C-5 methylcytosine is present as a 5mCpG
20 dinucleotide.

The invention further provides a method of treating a disorder of cell proliferation or development. The method involves administering to a subject a synthetic inhibitor molecule which recognizes and binds an allosteric site on DNA methyltransferase. The
25 binding of the synthetic inhibitor molecule prevents binding of AdoMet to DNA methyltransferase in a catalytically competent manner thereby inhibiting DNA methyltransferase. The inhibition of DNA methyltransferase prevents the methylation of DNA thereby treating the disorder of cell proliferation or development. In one embodiment, the synthetic oligonucleotide comprises a C-5 methylcytosine which
30 recognizes and binds an allosteric site on DCMTase thereby inhibiting DNA methyltransferase activity. In one embodiment, the disorder of cell proliferation is cancer, such as lung cancer or colon cancer. In one embodiment, the disorder of development is one linked to a genetic locus regulated by methylation. Examples of such disorders include, but are not limited to, Huntington's disease, Down's syndrome,
35 and disorders associated with a Hox gene.

The invention provides a method of inhibiting proliferation of cancer cells comprising administering to a subject a synthetic inhibitor molecule which recognizes and binds an allosteric site on DCMTase thereby resulting in an enzyme/synthetic inhibitor molecule complex, inhibiting DCMTase-mediated methylation of DNA, and thereby inhibiting proliferation of the cancer cells. In one embodiment, the cancer cell is from lung or colon. In one embodiment, the synthetic inhibitor molecule is an oligonucleotide comprising a C-5 methylcytosine which recognizes and binds an allosteric site on DCMTase thereby inhibiting DNA methyltransferase activity. In one embodiment, the C-5 methylcytosine is present as a 5mCpG dinucleotide. In one embodiment, the synthetic oligonucleotide comprises a nucleotide sequence as shown in Figure 1B and designated GC-box b^{MET} (SEQ ID NO:10), GC-box p^{MET} (SEQ ID NO:10), GC-box c^{MET} (SEQ ID NO:13), GC-box d^{MET} (SEQ ID NO:14), GC-box e^{MET} (SEQ ID NO:15), or CRE a^{MET} (SEQ ID NO:11). In one embodiment, the subject is a human. In another embodiment, the subject is an animal. In one embodiment, the animal is selected from a group consisting of porcine, piscine, avian, feline, equine, bovine, ovine, caprine and canine.

Definitions

All scientific and technical terms used in this application have meanings commonly used in the art unless otherwise specified. As used in this application, the following words or phrases have the meanings specified.

As used herein "synthetic oligonucleotide comprising a C-5 methylcytosine" means any non-naturally occurring oligonucleotide comprising a C-5 methylcytosine. The oligonucleotide can be a RNA, DNA or a derivative or hybrid thereof. The C-5 methylcytosine can be in the form of a 5mCpG dinucleotide. In one embodiment, the C-5 methylcytosine is centrally located within the oligonucleotide. In one embodiment, the synthetic oligonucleotide of the invention can be approximately 5 to approximately 70 bases in length. In another embodiment, the synthetic oligonucleotide can be approximately 15 to approximately 50 bases in length. In another embodiment, the synthetic oligonucleotide can be approximately 20 to approximately 30 bases in length. In another embodiment, the synthetic oligonucleotide is approximately 30 bases in length. Examples of synthetic oligonucleotides of the invention include, but are not limited to, the oligonucleotides GC-box b^{MET} (SEQ ID NO:10), GC-box p^{MET} (SEQ ID NO:10), GC-box c^{MET} (SEQ ID NO:13), GC-box d^{MET} (SEQ ID NO:14), GC-box e^{MET} (SEQ ID NO:15), and CRE a^{MET} (SEQ ID NO:11) shown in Figure 1B.

As used herein, "synthetic inhibitor molecule" includes synthetic molecules known in the art to facilitate entry of nucleic acids into cells and to minimize intracellular and intercellular breakdown of the nucleic acids. Examples of such antisense molecules
5 include, but are not limited to, peptide nucleic acid (PNA) and phosphorothioate-based molecules such as deoxyribonucleic guanidine (DNG) or ribonucleic guanidine (RNG). Also included are nonnucleic acid polymers derived from a library screen which bind the same site as the synthetic oligonucleotide of the invention.

10 As used herein, "an allosteric site" means a site other than an active site that can influence the catalytic progress of the enzyme. The influence can either inhibit or activate catalysis. For example, an active site on DCMTase includes the site to which AdoMet binds, the binding of AdoMet to the active site on DCMTase leading to the methylation of DNA. An active site is defined as the local protein environment in close
15 proximity to the reactive substituents in the methylation reaction.

As used herein, "DNA methyltransferase activity" means enzymatic activity that promotes transfer of a methyl group to DNA, thereby methylating DNA. An example of a source of a methyl group for transfer to DNA is AdoMet.

20

As used herein, "pharmaceutically acceptable salt" refers to a salt that retains the desired biological activity of the parent compound and does not impart any undesired toxicological effects. Examples of such salts include, but are not limited to, (a) acid addition salts formed with inorganic acids, for example hydrochloric acid,
25 hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; and salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, pamoic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acids, naphthalenedisulfonic acids, polygalacturonic acid; (b)
30 salts with polyvalent metal cations such as zinc, calcium, bismuth, barium, magnesium, aluminum, copper, cobalt, nickel, cadmium, and the like; or (c) salts formed with an organic cation formed from N,N'-dibenzylethylenediamine or ethylenediamine; or (d) combinations of (a) and (b) or (c), e.g., a zinc tannate salt; and the like. The preferred acid addition salts are the trifluoroacetate salt and the acetate
35 salt.

-16-

As used herein, "pharmaceutically acceptable carrier" includes any material which, when combined with a compound of the invention, allows the compound to retain biological activity and is non-reactive with the subject's immune system. Examples include, but are not limited to, any of the standard pharmaceutical carriers such as a phosphate buffered saline solution, water, emulsions such as oil/water emulsion, and various types of wetting agents. Preferred diluents for aerosol or parenteral administration are phosphate buffered saline or normal (0.9%) saline. Compositions comprising such carriers are formulated by well known conventional methods (see, for example, Remington's Pharmaceutical Sciences, Chapter 43, 14th Ed., Mack Publishing Co., Easton PA 18042, USA).

Compounds of the Invention

The invention provides a synthetic oligonucleotide comprising a C-5 methylcytosine and which recognizes and binds an allosteric site on DNA methyltransferase thereby inhibiting DNA methyltransferase activity. In one embodiment, the synthetic oligonucleotide has an inhibition constant of not greater than 1000 nM by steady-state kinetic assay. In another embodiment, the synthetic oligonucleotide has an inhibition constant of not greater than 200 nM by steady-state kinetic assay. In yet another embodiment, the synthetic oligonucleotide has an inhibition constant of not greater than 20 nM by steady-state kinetic assay. In one embodiment, the C-5 methylcytosine is centrally located within the oligonucleotide. In one embodiment, the synthetic oligonucleotide of the invention can be approximately 5 to approximately 70 bases in length. In another embodiment, the synthetic oligonucleotide can be approximately 15 to approximately 50 bases in length. In another embodiment, the synthetic oligonucleotide can be approximately 20 to approximately 30 bases in length. In a further embodiment, the synthetic oligonucleotide is approximately 30 bases in length. Examples of synthetic oligonucleotides of the invention include, but are not limited to, the oligonucleotides shown in Figure 1B and designated GC-box b^{MET} (SEQ ID NO:10), GC-box p^{MET} (SEQ ID NO:10), GC-box c^{MET} (SEQ ID NO:13), GC-box d^{MET} (SEQ ID NO:14), GC-box e^{MET} (SEQ ID NO:15), or CRE a^{MET} (SEQ ID NO:11).

Compositions Of The Invention

The invention further provides a composition comprising a synthetic oligonucleotide comprising a C-5 methylcytosine and which recognizes and binds an allosteric site on DNA methyltransferase. The composition is useful for inhibiting DNA

methyltransferase activity, thereby inhibiting the methylation of DNA. In one embodiment, the composition is a pharmaceutical composition comprising a pharmaceutically effective amount of a synthetic oligonucleotide comprising a C-5 methylcytosine, or a pharmaceutically acceptable salt thereof, and which recognizes and binds an allosteric site on DNA methyltransferase. In one embodiment, the pharmaceutical composition further comprises a pharmaceutically acceptable carrier. The pharmaceutical composition is useful for treating disorders associated with methylation defects, such as cancer and certain developmental disorders.

10 Administration of the Compositions

In accordance with the methods of the invention, the synthetic oligonucleotide can be administered in a pharmaceutical composition in unit dosage form. The most effective mode of administration and dosage regimen for the molecules of the present invention depend upon the location of the tissue or disease being treated, the severity and course of the medical disorder, the subject's health and response to treatment and the judgment of the treating physician. Accordingly, the dosages of the molecules should be titrated to the individual subject.

By way of example, the interrelationship of dosages for animals of various sizes and species and for humans based on mg/m^2 of surface area is described by Freireich, E.J., et al. Cancer Chemother., Rep. 50 (4): 219-244 (1966). It would be clear that the dose of the composition of the invention required to achieve an appropriate clinical outcome may be further reduced with schedule optimization.

25 Methods of the Invention

The invention further provides a method of inhibiting methylation of DNA. The method involves contacting a DCMTase with a synthetic inhibitor molecule in the presence of the DNA. The synthetic inhibitor molecule comprises a C-5 methylcytosine which recognizes and binds an allosteric site on DNA cytosine methyltransferase (DCMTase) thereby resulting in an enzyme/synthetic inhibitor molecule complex. The presence of the complex prevents DCMTase-mediated catalysis thereby inhibiting DCMTase activity and inhibiting methylation of DNA. In one embodiment, the synthetic oligonucleotide comprises a C-5 methylcytosine. In a further embodiment, the C-5 methylcytosine is present as a 5mCpG dinucleotide. Examples of synthetic inhibitor molecules include, but are not limited to, the oligonucleotides shown in Figure 1B and

designated GC-box b^{MET} (SEQ ID NO:10), GC-box p^{MET} (SEQ ID NO:10), GC-box c^{MET} (SEQ ID NO:13), GC-box d^{MET} (SEQ ID NO:14), GC-box e^{MET} (SEQ ID NO:15), or CRE a^{MET} (SEQ ID NO:11).

- 5 The invention further provides a method of treating a disorder of cell proliferation or development. The method involves administering to a subject a synthetic oligonucleotide which recognizes and binds an allosteric site on DCMTase. The binding of the synthetic oligonucleotide prevents DCMTase-mediated catalysis thereby inhibiting DCMTase activity. The inhibition of DCMTase prevents the methylation of
- 10 DNA thereby treating the disorder of cell proliferation or development. In one embodiment, the synthetic oligonucleotide comprises a C-5 methylcytosine. In one embodiment, the disorder of cell proliferation is cancer, such as lung cancer or colon cancer. In another embodiment, the disorder of development is one linked to a genetic locus regulated by methylation. Examples of such disorders include, but are not limited to,
- 15 to, Huntington's disease, Down's syndrome, and disorders associated with a Hox gene.

The invention provides a method of inhibiting proliferation of cancer cells comprising administering to a subject a synthetic inhibitor molecule which recognizes and binds an allosteric site on DCMTase thereby resulting in an enzyme/synthetic inhibitor molecule

20 complex. The presence of the complex prevents DCMTase catalysis thereby inhibiting DCMTase-mediated methylation of DNA, thereby inhibiting proliferation of the cancer cells. In one embodiment, the synthetic inhibitor molecule is an oligonucleotide comprising a C-5 methylcytosine which recognizes and binds an allosteric site on DCMTase thereby inhibiting DNA methyltransferase activity. In one embodiment, the

25 cancer is lung cancer or colon cancer. In one embodiment, the method of inhibiting proliferation of cancer cells comprises administering to a subject the synthetic oligonucleotide of the invention in a sufficient amount so that the oligonucleotide recognizes and binds an allosteric site on DCMTase so as to form an enzyme/synthetic oligonucleotide complex.

30 The invention provides a method of inhibiting hypermethylation of DNA comprising contacting a DNA cytosine methyltransferase (DCMTase) with a synthetic inhibitor molecule comprising a C-5 methylcytosine which recognizes and binds an allosteric site on DCMTase thereby resulting in an enzyme/synthetic inhibitor molecule complex, in

35 the presence of the DNA. The presence of the complex prevents DCMTase catalysis thereby inhibiting DCMTase activity and inhibiting hypermethylation of the DNA. In

one embodiment, the synthetic oligonucleotide comprises a C-5 methylcytosine. In a further embodiment, the C-5 methylcytosine is present as a 5mCpG dinucleotide. The inhibition of hypermethylation of DNA is useful for preventing the development of resistance to drugs such as anti-cancer drugs.

5

- The invention provides a method of inhibiting drug resistance in a subject comprising administering to a subject the synthetic oligonucleotide of the invention in a sufficient amount so that the oligonucleotide recognizes and binds an allosteric site on DCMTase so as to form an enzyme/synthetic oligonucleotide complex. The presence of the
- 10 complex prevents DCMTase catalysis so as to inhibit DCMTase-mediated hypermethylation of DNA thereby inhibiting drug resistance. The synthetic inhibitor molecule can be administered to a subject prior to, concurrent with or after administration of an anti-cancer therapeutic agent to prevent overmethylation of DNA induced in the subject's cells in response to the anti-cancer therapeutic agent.
- 15 The invention additionally provides a method for screening molecules, such as those obtained from a combinatorial library, to identify modulators of DCMTase which recognize and bind an allosteric site on DCMTase. In one embodiment, the modulator is an inhibitor of DCMTase. In another embodiment, the modulator is an activator of DCMTase. The method comprises contacting a molecule with DCMTase in the
- 20 presence of AdoMet and DNA, and measuring DCMTase activity. An increase in DCMTase activity is indicative of an activator of DCMTase and a decrease in DCMTase activity is indicative of an inhibitor of DCMTase. DCMTase activity can be measured by methods known in the art, including the assays disclosed in the Examples provided herein. Those of ordinary skill in the art can identify a modulation
- 25 of DCMTase activity that is indicative of binding an allosteric site on the enzyme. In a preferred embodiment, DCMTase activity is measured by a steady-state assay. One can plot enzyme activity as a function of varied concentrations of the molecule being tested, and also as a function of varied concentrations of DNA and AdoMet. Preferably, a mathematical fit is performed on the plotted results. Competitive
- 30 inhibition by DNA and uncompetitive inhibition by AdoMet, or competitive inhibition by AdoMet and uncompetitive inhibition by DNA, for example, would be indicative of an inhibitor molecule which recognizes and binds an allosteric site on DCMTase. Also included within the invention are modulators of DCMTase which recognize and bind an allosteric site on DCMTase, and which are identified by the above method.

Advantages of the Invention

The invention disclosed herein provides a potent and reversible inhibitor of DNA methyltransferase that does not require incorporation into DNA. This inhibitor can be used to inhibit methylation of DNA and to treat disorders associated with DNA methylation defects, such as cancer and developmental disorders.

In addition to identifying particular synthetic oligonucleotides which inhibit DNA methyltransferase, the invention provides information about the mechanism responsible for this inhibition. By identifying an allosteric site on DCMTase as the site of action of the inhibitors, the invention provides a basis for developing and identifying variants of the particular synthetic oligonucleotides disclosed herein that will also be useful for inhibiting DNA methyltransferase. Additionally, the disclosure herein teaches that a C-5 methylcytosine is responsible for the potency of the inhibition effected by the synthetic oligonucleotides of the invention.

EXAMPLES

The following examples are presented to illustrate the present invention and to assist one of ordinary skill in making and using the same. The examples are not intended in any way to otherwise limit the scope of the invention.

Example 1: DNA Binding Discrimination of the Murine DNA Cytosine-C⁵ Methyltransferase

In this example gel mobility shift analyses (GMSA) using defined sequences to estimate K_D^{DNA} and *in vitro* screening method of a large, divergent pool of DNA, are used to determine discrimination of DCMTase. The results presented herein demonstrate that the DCMTase:DNA complex is concluded to be thermodynamically stabilized by guanosine/cytosine-rich sequences flanking a central CpG cognate site.

Materials

DCMTase was purified from mouse erythroleukemia cells as previously described (Xu, G., et al. (1995) Biochemi. Biophys. Res. Communi. 207:544-551). S-adenosyl-L-[methyl-³H]methionine (75 Ci/mmol, 1 mCi/ml, 1 Ci=37 GBq) was from Amersham Life Sciences (Arlington Heights, Illinois), Unlabeled AdoMet, purchased from Sigma Chemical Company (St. Louis, MO), was further purified as

described (Reich, N.O. & Mashhoon, N. (1990) Inhibition of *EcoRI* DNA methylase with cofactor analogs. J. Biol. Chem. 265:8966–8970). Routinely, a 125 mM AdoMet stock concentration was prepared at a specific activity of 5.8×10^3 cpm/pmol. DE81 filters were purchased from Whatman Inc. (Lexington, MA). All other chemicals and reagents were purchased from Sigma Chemical Company (St. Louis, MO) or Fisher Scientific (Hampton, New Hampshire).

DNA Substrate Preparation

The preparation, purification, and analysis of six oligonucleotides that mimic the GC-box and the cyclic AMP responsive elements (CRE) were previously described (Flynn, J., et al., 1996, Murine DNA cytosine–C5 methyltransferase: Pre-steady and steady-state kinetic analyses with regulatory DNA sequences, Biochemistry 35:7308–7315) (Figure 1). The percentage of double-stranded DNA in annealed DNA samples was confirmed to be greater than 99% by ^{32}P -radiolabeling, polyacrylamide gel separation, subsequent autoradiography and densitometry using a CCD camera and the SW5000 analysis package from Ultra Violet Products (UVP, San Gabriel, CA).

Gel Mobility Shift Assays

Gel mobility shift assays (GMSA) were performed with minor revisions to the original procedures (Fried, M., & Crothers, D.M., 1981, Equilibria and kinetics of the lac repressor–operator interactions by polyacrylamide gel electrophoresis, Nucleic Acids Res. 9:6505–6525; Garner, M.M. & Revzin, A., 1981, A gel electrophoresis method for quantifying the binding proteins to specific DNA regions: applications to components of the Escherichia coli lactose operon regulatory system, Nucleic Acids Res. 13:3047–3060). All reactions were done in 100 mM Hepes pH 7.4, 10 mM EDTA, 10 mM DTT, 200 mg/ml BSA, 5% glycerol using the indicated ^{32}P -labeled DNA and DCMTase concentrations, incubated on ice for 5 minutes and loaded on a 1xTBE (89 mM Tris–HCl pH 8.3, 89 mM boric acid, 2 mM EDTA), 6% polyacrylamide gel. Electrophoresis was done at 250 V, 9 mA for 2 hours at 4 °C and the dried gel was exposed to film overnight. The reaction conditions for buffer, temperature, incubation time, cofactor addition and gel composition have all been optimized. Only slightly better complex resolution was obtained under the listed conditions compared to a 10 minute incubation at 37 °C prior to gel loading at room temperature and containing either cofactor S-adenosyl–L-methionine, product S-adenosyl–L-homocysteine, or the AdoMet analog sinefungin. Hepes reaction buffer at

pH 7.4 produced sharper banding than Tris-HCl at pH 8.0. Initial binding assays, with a limiting, and constant DNA concentration, resulted in the formation of multiple bands. Subsequent assays used a limiting and constant enzyme concentration with varying DNA concentrations.

5

Binding Isotherm Determinations of K_D^{DNA}

Autoradiogram-derived band intensities corresponding to the mobility shifted DCMTase:DNA complexes were acquired using the UVP system described above. Background subtractions were from equivalent areas about one centimeter below each mobility shifted complex. The corrected intensities were then fit to a nonlinear binding isotherm and graphed using KaleidaGraph 2.1.2 software (Synergy Software, Reading, PA) The intensity of the labeled DNA in the protein:DNA complex at saturation was directly compared to uncomplexed DNA areas in control lanes containing 50%, 100% and 150% molar DNA equivalents of the DCMTase concentration.

15

Screening for DNA binding preferences

An *in vitro* selection approach was used to determine the DNA binding discrimination of DCMTase. A population of DNA molecules, each 66 base pairs long, were synthesized with a central CpG dinucleotide flanked on each side by 12 positions randomized with either adenosine, thymidine or cytidine; total complexity equal to 2.8×10^{11} discrete sequences (Figure 6). Guanosine was not added to the randomization to avoid multiple CpG dinucleotides on a double-stranded DNA. The randomized regions are flanked by PCR primer regions that contain the restriction sites used for cloning. The first generation pool of DNA was made double-stranded by Klenow polymerase extension of primer D.

20

25

The screening procedure was reiterated five times under the conditions listed in Table 2 (see Results, *infra*). DNA substrates from each pooled generation that induced higher thermodynamic stabilities of the DCMTase:DNA complex were separated from lower affinity DNA by PAGE as described above. The region of the gel containing shifted DNA complexes was excised and five exchanges of 5 mL water over 72 hours shaking on ice was sufficient to elute greater than 95% of all cpm present in the excised gel slice as determined by Cerenkov counting. The eluted DNA was lyophilized, resuspended in TE (10 mM Tris-HCl at pH 8.0; 1 mM EDTA) and

30

35

cleaned by one phenol:chloroform and two chloroform extractions followed by ethanol precipitation and resuspension in TE. The selected DNA pools were amplified using 20 rounds of PCR using Deep Vent polymerase (New England Biolabs) and the DNA primers shown in Figure 6. The 66 base pair DNA was separated from the PCR
5 primers on agarose gels and purified using minor changes to the original procedure (Wieslander, L., 1979, A simple method to recover intact high molecular weight RNA and DNA after electrophoretic separation in low gelling temperature agarose gels, Anal. Biochem. 98:305-3).

10 *Identification of preferred DNA substrates*

Individual members from the selected DNA pools were identified by cleaving the DNA ends with *Bam*HI and *Eco*RI endonuclease and cloning into pGEM11zf- (Promega, Madison, WI) using standard protocols. The plasmid DNA from single isolates was prepared and the selected CpG flanking sequences were determined using
15 the CircumVent sequencing kit (New England Biolabs, Beverly, Massachusetts). The selected inserts were sequenced from both strands using the T7 and SP6 sequencing primers (Promega, Madison, WI). Statistical analyses were performed using several programs in the Wisconsin Sequence Analysis Package (Genetics Computer Group, Madison, WI) and Kaliedagraph (Synergy Software, Reading, PA). Statistical
20 significance was determined by the Student's t-Test using Microsoft Excel, Microsoft, Redmond, WA.

The selected generations were analyzed for initial velocity. The 50 mL reactions contained 50 nM DCMTase, 7 mM AdoMet and DNA at 4.7, 23, 47 and 230 nM in
25 100 mM Tris pH 8.0, 10 mM EDTA, 10 mM DTT, 200 mg/mL BSA. AdoMet (*S*-adenosyl-L-[methyl-³H]methionine ([methyl-³H]AdoMet) (75 Ci/mmol, 1mCi/ml, 1 Ci = 37 Gbq) was purchased from Amersham (Arlington Heights, IL). The incubations were for 1 hour at 37 °C.

30 DNA with tritiated C-5 cytosines, deposited by the DCMTase, were separated from the tritiated AdoMet by spotting the reaction on DE 81 filters (Whatman, Lexington, MA) followed by a series of 200 mL washes; three in 50 mM HK₂PO₄ and one each in 80% ethanol, 95% ethanol and ethyl ether. Dried filters were placed in 3 mL of LiquiScint (National Diagnostics, Atlanta, GA) and counted in a scintillation counter.
35 Counts per minute were transformed to femtomoles of methyl groups deposited on DNA over the course of the reaction.

Results

Gel mobility shift analyses of GC-box and CRE cis-elements

The preliminary experiments used a standard gel mobility shift assay in which a constant, low DNA concentration was titrated with higher protein concentrations. As shown in Figure 2, essentially all of the GC-box a/b (100 pM) binding occurred between 5 nM and 95 nM DCMTase. An initial complex was formed at the lower DCMTase concentrations and an abrupt shift of most of the free DNA is coincident with the formation of a second complex at about 20 nM DCMTase. Further addition of DCMTase resulted in the loss of the more mobile complex I in favor of a less mobile complex II. Similar results were observed for GC-box a/b^{MET}, CRE a/b, and CRE a^{MET}/b. The complexes shown in Figure 2 contained the DCMTase, as addition of an antibody to DCMTase resulted in an increased shift of each complex. Coincubation of DCMTase and GC-box a/b with a 40-fold excess of unlabeled polydA:polydT, calculated on a dinucleotide basis, did not disrupt the specific DCMTase:DNA complex.

The multiple banding of DCMTase:DNA complexes observed in Figure 2 are similar to results obtained with two cytosine DNA methyltransferases, *M.MspI* (Dubey, A.K. & Roberts, R.J., 1992, Sequence-specific DNA binding by the *MspI* DNA methyltransferase, *Nucleic Acids Res.* 20:3167-3173) and *M.HhaI* (Mi, S. & Roberts, R.J., 1993, The DNA binding affinity of *HhaI* methylase is increased by a single amino acid substitution in the catalytic center, *Nucleic Acids Res.* 21:2459-2464; Reale et al., 1995, DNA binding and methyl transfer catalyzed by mouse DNA methyltransferase, *Biochem. J.* 312:855-861) obtained similar gel shift results with a mammalian DCMTase and assumed that the slower migrating band contained two DCMTase molecules bound to a single DNA.

Steady-state kinetic analyses of the DCMTase with the same 30 base-pair DNA substrates used in these studies indicate that K_m^{DNA} is 1000- to 50,000-fold higher (Flynn, J., et al., 1996, Murine DNA cytosine-C5 methyltransferase: Pre-steady- and steady-state kinetic analyses with regulatory DNA sequences, *Biochemistry* 35:7308-7315) than the DNA concentrations used to generate Figure 2 and used by Reale et al, 1995, *supra*. The complexes formed in Figure 2 under limiting DNA and excess protein do not promote a detectable catalytic activity. Therefore, the stability of

protein:DNA was determined by keeping the enzyme at a constant concentration (100 nM) and varying the amount of added DNA (Dubey & Roberts, 1992, *supra*). Figures 3 through 5 show the results with this approach for GC-box a/b^{MET}, GC-box a/b and GC-box b^{MET}. In all cases a single shifted band is resolved, the binding isotherm data are fit well by a simple hyperbola, and each complex is saturable. The apparent K_D^{DNA} estimations for the different forms of GC-box and CRE are summarized in Table 1. The formation of equimolar protein:DNA complexes is supported by comparisons of the band intensity for complexed DNA at saturation with the band intensities of control reactions containing 50, 100 and 150 nM DNA and no enzyme (Figures 3 and 4, lanes 1, 2 and 3).

Table 1: Determinations of K_D^{DNA} for DCMTase by Gel Mobility Shift Assay.^a

	DNA Substrate	K_D^{DNA} (nM)
15	GC-box a	1.2 +/- 0.2
	GC-box b	1.3 +/- 0.4
	GC-box b ^{MET}	0.88 +/- 0.13
	GC-box a/b	0.42 +/- 0.15
20	GC-box a/b ^{MET}	0.36 +/- 0.11
	CRE a	>50
	CRE b	>50
	CRE a/b	1.5 +/- 0.4
25	CRE a ^{MET} /b	1.0 +/- 0.3

^a The values presented were obtained from the relative intensities of bands corresponding to DCMTase:DNA complexes fit by non-linear regression as described in Experimental Procedures.

30

DCMTase:DNA complexes formed by high substrate concentrations travel with the same relative mobility as complex I in Figure 2. For DNA concentrations higher than about 10 times the apparent K_D^{DNA} , the complexes become less mobile. The complex formed between DCMTase and single-stranded GC-box b^{MET} (Figure 5, lane 7) is shown to migrate to approximately the same distance as the complex formed between DCMTase and hemi-methylated CRE a^{MET}/b DNA (lane 9).

35

The estimates of apparent K_D^{DNA} are consistent with our previous K_m^{DNA} estimates (Table 1 and Flynn, J., et al., 1996, Murine DNA cytosine-C5 methyltransferase: Pre-steady- and steady-state kinetic analyses with regulatory DNA sequences, Biochemistry 35:7308-7315). The hemi-methylated double-stranded form of DNA was bound with a slightly higher affinity than the unmethylated double-stranded form for both the GC-box and CRE DNA. In support of CpG flanking sequence discrimination by DCMTase, the GC-box substrates of each duplex form had an approximate three-fold lower K_D^{DNA} than the corresponding CRE DNA form. Single-stranded substrates bound with less stability than double-stranded DNA. The binding of CRE single-strands was exceptionally poor and at the limits of resolution by this technique. GMSA was capable of resolving a binding discrimination in favor of guanosine/cytosine-rich sequences flanking a central CpG dideoxynucleotide.

15 *Screening for DCMTase binding discrimination with a randomized DNA pool*

The sampling of several discrete sequences for binding specificity is laborious and prone to investigative prejudice. In order to understand the thermodynamic stability of DCMTase:DNA interactions in a diverse population of CpG sequence contexts, as might be expected *in vivo*, we devised an *in vitro* screening protocol that exploits the gel mobility shift assay. The reaction conditions used for each iterative generation of the screening are summarized in Table 2. The first round of screening contained ten times more DNA molecules than the maximal population complexity of 2.8×10^{11} discrete sequences. An increasing fraction of the added randomized pool was shifted through the first three generations, during which the enzyme concentration was kept constant and the DNA concentration was decreased. The initial conditions were sufficient to stabilize binding of the DNA pool, so the selective pressure to discriminate between sequences was increased in generation-4 and 5 by decreasing both enzyme and DNA concentrations. The maximal population complexity in each generation decreases because only a fraction of the added DNA was shifted. The complexity of the starting population is divided by the percentage of DNA shifted in each generation and ultimately results in no more than 1.2×10^4 discrete sequences in the generation-5 pool (Table 2).

Table 2: Binding conditions and gel shift results of *in vitro* screening^a

	Iterative	DCMTase	DNA	Percentage	Maximal
5	Generation	Concentration	Concentration	*DNA Shifted	Population Complexity
	0	—	—	—	2.8x10 ¹¹
	1	68 nM	50 nM	1.5%	4.2x10 ⁹
	2	68 nM	25 nM	12%	5.0x10 ⁹
10	3	68 nM	12.5 nM	25%	1.2x10 ⁸
	4	5.0 nM	0.125 nM	<1%	1.2x10 ⁶
	5	0.50 nM	0.030 nM	<1%	1.2x10 ⁴

^a Listed are the enzyme and DNA substrate concentrations used in each round of selection. The Cerenkov cpm within the excised gel slice, containing the shifted complex, is shown as a percentage of the total Cerenkov counts loaded onto the gel. This percentage limits the complexity of the DNA pool, therefore it is used to calculate the maximal population complexity in each successive generation.

Individual members from the starting pool and generations—1, 3 and 5 were cloned and sequenced from both strands. Only the guanine containing strands are shown for simplicity in Figure 7, however, these studies were done using unmethylated double-stranded substrates. Synthesis of the starting population is shown to be randomized at each position with the expected frequency approximating 1/3 each in guanine, adenine and thymine.

The selected pools successively became more guanosine-rich with each generation. A total of 49 isolates were cloned and sequenced from the generation—5 pool and none were identical. Nucleotide, dinucleotide and trinucleotide frequencies were analyzed using the COMPOSITION (Wisconsin Sequence Analysis, Madison, WI). The selected nucleotides flanking the central CpG dinucleotide were 64.7% in guanine, 13.8% in adenine and 21.6% in thymine. The mean frequency of guanine bases per generation—5 isolate was 14.5 out of the 24 selectable positions and more guanines were observed on the 5'-flank compared to the 3'-flank ($p = 0.04$). The far flanking regions are a full helical turn distal to the invariant CpG and are highly enriched in guanine as compared to regions proximal to the CpG. In addition to the abundance of

guanosyl-guanosyl (GpG) dinucleotides, guanosyl-thymidyl (GpT) and thymidyl-guanosyl (TpG) dinucleotides appear often and occur more frequently on the 3'-flank ($p = 0.01$). Trinucleotide analyses reinforce the observations at the nucleotide and dinucleotide levels. The highest frequency of GpGpG was at the far 5'-flank, while
5 GpTpG and TpGpT trinucleotides were far more abundant in the 3' flank ($p = 0.01$). The discrimination exhibited by DCMTase for generation-5 sequences may reflect an important structural characteristic that contributes to stabilization of the initial DCMTase:DNA complex. These analyses suggest that an ideal substrate has sequence asymmetry around the CpG and that there is a particular binding orientation of
10 DCMTase on DNA.

The guanine-richness at each randomized position for the generation-5 isolates is best shown in Figure 8. The murine DCMTase is a large 183,000 Da protein that selected for sequences extending over the entire 12 base-pairs provided for selection
15 on each side of the central CpG. The Wisconsin Sequence Analysis program CONSENSUS was used to construct a common generation-5 sequence with a certainty level of 60%. The sequence GGGGGGRRKKGCGKGGKGGKGGKGG (SEQ ID NO:1), where R is guanine or adenine and K is guanine or thymine, was obtained and is shown to highlight the guanine richness and the preference for GpT and TpG on the 3'-side of the CpG. At a certainty level of 80% the plasticity of
20 sequence preferences can be seen close to the invariant CpG; KGGRRKRRDDKRCGKRRDKKKKKKKKG (SEQ ID NO:2) (D is guanine, thymine or adenine). We have not tested whether the DCMTase can select for sequences out further than 12 base-pairs or if multiple CpG dinucleotides are preferred over the 26
25 base-pair expanse.

Similar sequences occur frequently in the genome

We subjected the 49 generation-5 sequences to FASTA searches of the GenBank library to see if similar sequences exist in the genomes of higher eukaryotes. The
30 search was limited in three ways. First, only the mouse and human sequences were searched, even though DCMTase activities have been identified in many metazoan organisms. Second, to be considered further, a "hit" had to be identical at 22 of the 26 base positions, including the central CpG. No hits were retrieved that had a higher identity. Third, no gaps in alignment were allowed.

Remarkably, 20 "hits" were recovered from GenBank that met these severely restricted criteria. Figure 9 shows the alignments of the five hits from mouse and lists the 15 hits from human. A simplified, random genome would be expected have a complexity of 4^{22} , or 1.8×10^{13} base-pairs, in order to contain any of these sequences just once. Of course, this is an oversimplification. But, the results appear to be striking when considering the mammalian genome is approximately 3×10^9 base-pairs, only about 40% in guanine plus cytosine, and about 10-fold deficient in CpG dinucleotides. The majority of hits are in what may be presumed to be regulatory regions of the genome; 5' or 3' untranslated regions (UTR) or in CpG islands. Many of the associated genes are also of developmental interest. For example, homeo box Hox2.6 and HoxA7 function in early body segmentation. These findings may reflect an intrinsic function of DCMTase in developmental programming.

Control experiments eliminate a non-specific selection

A control series of amplifications in the absence of DCMTase were done to show that our iterative PCR conditions were not responsible for the guanine selection observed with the generation-5 DNA. Endonuclease challenge was done with *Taq* I (5'-TCGA-3' restriction) and *Aci* I (5'-GCGG-3' restriction) to assess the randomness of mock selected and DCMTase selected pools. Although this is a limited sampling, the DNA specificity of these enzymes can discern the relative abundance of nucleotides immediately flanking the CpG. The guanine-richness is probed by *Aci* I and the adenine/thymine-richness is probed by *Taq* I. After endonuclease challenge of ^{32}P labeled DNA, the products were resolved on a 12% polyacrylamide gel. Using densitometry, the intensities of the restricted bands were compared to unrestricted control bands. The mock generation-5 sequences immediately flanking the CpG remained random, approximately 6% of the DNA was restricted by each endonuclease, demonstrating that a non-specific selection did not occur under these experimental conditions. Results with the DCMTase selected generations were consistent with the guanine/cytosine selection determined from sequencing individual clones. Also, consistent with a lack of selection from the protocols alone, the entire mock selected pool was sequenced and compared to the DCMTase selected generation-5 pool, similar to that done by Blackwell et al., 1991. An equal abundance of the randomized nucleotides was resolved for the mock selected pool and a guanine-rich population was resolved for the DCMTase selection.

The DCMTase-selected DNA from the iterative generations were compared to each other in binding and catalytic assays. The DCMTase binds the pooled generation-5 sequences only two-fold more tightly than the starting pool. The inherent complexity of each pool makes it difficult to assess the true preference for each generation as a whole. The question of sequence specificity was more accurately addressed by GMSA
5 of the discrete sequences, CRE a/b and GC-box a/b. There we found that the guanine/cytosine-rich GC-box was preferred approximately 3-fold compared to the more adenine/thymine-rich CRE sequence. Figure 10 shows the initial velocity plots for the starting population and generations-2, 4 and 5. The catalytic specificity for the selected generations increases at each cycle, with little change in K_m^{DNA} and a two-
10 fold increase in k_{cat} .

Discussion

Because it is the catalytic agent for cytosine methylation, DCMTase clearly has a central role in both maintaining DNA methylation patterns and in establishing new
15 "epi-genotypes". The fundamental issues of binding and catalytic discrimination of the mammalian enzyme for different DNA sequences have been actively debated. Many reports have suggested that the ability of the enzyme to methylate the cognate CpG dinucleotide depends to some degree on flanking sequences (Bolden, A.H., et al.,
20 1986, Primary DNA sequence determines sites of maintenance and de novo methylation by mammalian DNA methyltransferases, Mol. Cell. Bio. 6:1135-1140; Bestor, T.H., et al., 1992, CpG islands in mammalian gene promoters are inherently resistant to de novo methylation, GATA 9:48-53; Hepburn, P.A., et al., 1991, Enzymatic methylation of cytosine in DNA is prevented by adjacent O⁶-
25 methylguanine residues, J. Biol. Chem. 266:7985-7987; Pfeifer, G.P., et al., 1985, Mouse DNA-cytosine-5-methyltransferase: sequence specificity of the methylation reaction and electron microscopy of enzyme-DNA complexes, EMBO J. 4:2879-2884; Ward, C., et al., 1987, *In vitro* methylation of the 5'-flanking regions of the mouse b-globin gene, J. Biol. Chem. 262:11057-1106; Carotti, D., et al., 1986,
30 Substrate preferences of the human placental DNA methyltransferase investigated with synthetic polydeoxynucleotides, Biochim. et Biophys. Acta 866:135-143; Smith, S.S., et al., 1992, Mechanism of human methyl-directed DNA methyltransferase and the fidelity of cytosine methylation, Proc. Natl. Acad. Sci. USA 89:4744-4748), while others describe the lack of any flanking sequence effects (Bestor, T.H. and Tycko, B.,
35 1996, Creation of methylation patterns, Nature Genetics 12:363-367; Carlson, L., et al., 1992, Properties and localization of DNA methyltransferase in preimplantation

embryos: implications for genomic imprinting, *Genes and Development* 6:2536–2541). These studies used partially purified or proteolyzed enzyme, substrates containing multiple CpG sites, and compared relative velocities obtained at a single substrate DNA concentration, thereby precluding an accurate estimation of specificity
 5 (otherwise known as discrimination).

Similarly, reports regarding the preference of DCMTase for single- and double-stranded substrates are also in direct conflict with one another (Adams, R.L.P., et al., 1986, Mouse ascites DNA methyltransferase: characteristic of size, proteolytic
 10 breakdown and nucleotide recognition, *Biochim. Et Biophys. Acta* 868:9–16; Smith et al., 1992, *supra*; Carotti et al., 1986, *supra*; Wang, R.Y.H., et al., 1984, Human placental DNA methyltransferase: DNA substrate and DNA binding specificity, *Nucl. Acids Res.* 12:3473–3490; Pfeifer et al., 1985, *supra*; Gruenbaum, Y., et al., 1982, Substrate and sequence specificity of a eukaryotic DNA methylase, *Nature* 295:620–
 15 622; Christman, J.K., et al., 1995, 5-Methyl-2'-deoxycytidine in single-stranded DNA can act in *cis* to signal *de novo* DNA methylation. *Proc. Natl. Acad. Sci. USA* 92:7347–7351).

A recent steady-state kinetic analysis with unmethylated GC-box and CRE DNA
 20 sequences showed compensatory 3- to 4-fold changes in K_m^{DNA} and k_{cat} that resulted in a small discrimination at the level of k_{cat}/K_m^{DNA} (Flynn, J., et al., 1996, Murine DNA cytosine-C5 methyltransferase: Pre-steady- and steady-state kinetic analyses with regulatory DNA sequences, *Biochemistry* 35:7308–7315). In this Example, the sequence-dependent discrimination of DCMTase is quantitatively addressed at the
 25 level of K_D^{DNA} . The thermodynamic binding constant, K_D^{DNA} , is a characteristic of the initial enzyme:DNA complex and K_m^{DNA} has an additional term accounting for the forward reaction rate. DCMTase:DNA interactions were investigated with discrete DNA sequences of biological importance, and with a large divergent pool of DNA sequences. The discrimination between unmethylated single- and double-stranded
 30 DNA, and unmethylated and hemi-methylated double-stranded DNA was also quantified.

DCMTase binding to DNA is stabilized by guanine/cytosine-rich sequences

Gel mobility shift assays were used to determine the apparent dissociation constants,
 35 K_D^{DNA} , of the enzyme for different forms of the GC-box and CRE *cis*-regulatory elements. Complex, higher-order interactions were observed under the more standard

conditions of limiting DNA and varying protein concentrations. While the multiple protein:DNA complexes and unusual DNA concentration dependence are shown to involve the DCMTase, accurate quantitative analysis is precluded due to the uncertainty of binding stoichiometry and the relative affinities of each binding event

5 (Senear, D.F., & Brenowitz, M., 1991, Determination of binding constants for cooperative site-specific protein-DNA interactions using the gel mobility shift assay, J. Biol. Chem. 266:13661-13671; Sackett, D.L. & Saroff, H.A., 1996, The multiple origins of cooperativity in binding to multi-site lattices, FEBS 397:1-6). Whereas many DNA-binding proteins, including DNA adenine-N⁶ methyltransferases (Reich,

10 N.O. & Mashhoon, N., 1990, Inhibition of *EcoRI* DNA methylase with cofactor analogs, J. Biol. Chem. 265:8966-8970), form a single protein:DNA complex under similar conditions, bacterial and mammalian DNA cytosine C⁵ methyltransferases are known to produce multiple complexes at low DNA concentrations (Dubey, A.K. & Roberts, R.J., 1992, Sequence-specific DNA binding by the *MspI* DNA

15 methyltransferase, Nucleic Acids Res. 20:3167-3173; Mi, S. & Roberts, R.J., 1993, The DNA binding affinity of *HhaI* methylase is increased by a single amino acid substitution in the catalytic center, Nucleic Acids Res. 21:2459-2464; Reale, A., et al., 1995, DNA binding and methyl transfer catalyzed by mouse DNA

20 methyltransferase, J. Biochem. 312:855-861). The multiple complexes formed with excess enzyme and DNA concentrations far below K_m^{DNA} may be common to cytosine DNA methyltransferases. These complexes are known to be catalytically incompetent in the case of the murine enzyme (Flynn, J., et al., 1996, Murine DNA cytosine-C5 methyltransferase: Pre-steady- and steady-state kinetic analyses with regulatory DNA sequences, Biochemistry 35:7308-7315).

25

Gel mobility shift assays performed with micromolar DNA concentrations and limiting DCMTase result in a single, shifted DNA band. These observations are again similar to those described for the bacterial cytosine DNA methyltransferases, *M.MspI* (Dubey & Roberts, 1992, *supra*) and *M.HhaI* (Mi & Roberts, 1993, *supra*); the

30 determination of equilibrium constants under these conditions is valid and not. In fact, our enzyme preparation obeyed classical Michaelis-Menton kinetics with the same substrates when assayed in the same DNA concentration range (Flynn et al., 1996, *supra*). Also, the estimated K_D^{DNA} values reported in Table 1 are similar to those previously reported at the level of K_m^{DNA} with the same DNA (Flynn et al., 1996,

35 *supra*). The K_D^{DNA} values are about one-half of those determined at the level of K_m^{DNA} for the same double-stranded substrates. The lack of large differences between

these constants suggests that steps following the initial formation of a specific protein:DNA complex do not contribute largely to K_m^{DNA} .

DCMTase bound DNA in a 1:1 stoichiometry and had a strong preference for binding
 5 double-stranded DNA over single-stranded DNA. Hemi-methylated DNA was
 bound by the enzyme with slightly higher affinity than unmethylated double-stranded
 DNA. The K_D^{DNA} data further supports the interpretation that the preference for hemi-
 methylated DNA versus unmethylated double-stranded DNA derives almost entirely
 from changes in the methylation rate constant, $k_{\text{methylation}}$ (Flynn, J., et al., 1996,
 10 Murine DNA cytosine-C5 methyltransferase: Pre-steady- and steady-state kinetic
 analyses with regulatory DNA sequences, *Biochemistry* 35:7308-7315). A recent
 study of *M.HhaI*:hemi-methylated DNA and *M.HhaI*:DNA cocrystal structures
 attempted to rationalize the two to three-fold discrimination manifested by this
 enzyme at the level of binding (O'Gara M., et al., 1996, A structural basis for the
 15 preferential binding of hemimethylated DNA by *HhaI* DNA methyltransferase, *J. Mol.*
Bio. 263:597-606). These authors proposed that the binding discrimination derives
 mostly from a single van der Waals' contact between the Glu²³⁹ carboxylate and the
 methyl group of the 5-methyl-2'deoxyctidine. While the DCMTase also has a
 glutamate at this position (Glu¹³⁸⁸), we suggest that other differences in the assembly
 20 of the active site contribute to the quantitatively larger preference for hemi-
 methylated DNA shown by the murine enzyme.

The two base-pair, CpG, cognate sequence of the mammalian DCMTase is small
 compared to the cognate sites of most bacterial DNA methyltransferases. DNA
 25 footprint analyses of *M.SssI*, *M.HhaI* and *M.MspI* are consistent with protein:DNA
 interactions extending over 16 base-pairs (Renbaum, P. & Razin, A., 1995, Footprint
 analysis of *M.SssI* and *M.HhaI* methyltransferases reveals extensive interactions with
 the substrate DNA backbone *J. Mol. Biol.* 248:19-26; Dubey, A.K. & Roberts, R.J.,
 1992, Sequence-specific DNA binding by the *MspI* DNA methyltransferase, *Nucleic*
 30 *Acids Res.* 20:3167-3173). Thus, the large mammalian DCMTase protein (Glickman,
 J.F. & Reich, N.O., 1997, Peptide mapping of the murine DNA methyltransferase
 reveals a major phosphorylation site and the start of translation, *J. Biol. Chem.* In
 press) most likely involves DNA contacts outside of this minimal sequence. Support
 for this is provided by the observation that the guanine/cytosine-rich GC-box element
 35 (GGGGCGGGGC (SEQ ID NO:3)) is bound approximately 3-fold more tightly than
 the adenine/thymine-rich CRE element (TGACGTCA). An *in vitro* selection method

was designed to define both the span of the protein:DNA interface, and the sequence preference of the enzyme for nucleotides flanking the consensus CpG. Previous applications of this strategy were useful in defining a consensus sequence for DNA binding proteins involving large differences in binding energetics between random and target sequences (Kinzler, K.W. & Vogelstein, B., 1989, Whole genome PCR: application to the identification of sequences bound by regulatory proteins, Nucl. Acids Res. 17:3645-3653; Thiesen, H. & Bach, C., 1990, Target detection assay (TDA): A versatile procedure to determine DNA binding sites as demonstrated on SP1 protein, Nucl. Acids Res. 18:3203-3209; Blackwell, T.K., et al., 1990, Sequence-specific binding by the c-Myc protein, Mol. Cell. Bio. 13:5216-5224; He, Y., Stockley, P.G. & Gold, L., 1996, *In vitro* evolution of the DNA binding sites of *Escherichia coli* methionine repressor, MetJ. J. Mol. Biol. 255:55-66). These selection strategies were extended to identify flanking sequence preferences, where binding discrimination is expected to be much less than when searching for a six to ten base-pair cognate site. One potential outcome would be the lack of any preference, as described for the UBF protein using this method (Copenhaver, G.P., et al., 1994, The RNA polymerase I transcription factor UBF is a sequence-tolerant HMG-box protein that can recognize structured nucleic acids, Nucleic Acids Research 22:2651-2657). A consensus sequence larger than the minimal CpG was not likely to result from this selection process, because genomic sequencing of 5-^mC reveals that the enzyme methylates many CpG contexts *in vivo*.

The screening method employed herein efficiently identified a DCMTase-induced population drift from 33.3% guanosine in the starting randomization to 50.0% in generation-1, 55.3% in generation-3 and finally 64.7% in generation-5. Randomized position 12 (see Figure 8) was enriched to 88% guanine in generation-5, suggesting that the total sequence space represented by the starting randomization was severely confined. Ultimately, the selection process did not disclose an obvious preferred sequence, but clearly a selection was evident. This is consistent with the observation that roughly 3×10^7 CpG flanking sequence contexts in the murine genome undergo methylation *in vivo*.

Sequence analysis of the 49 generation-5 members provided evidence that the DCMTase may bind these substrates in a preferred orientation. A greater guanosine selectivity was associated with the far 5'-side of the CpG and a more divergent region was exposed from the -2 to the -5 positions. The 3'-side of the invariant CpG

exhibits a different DCMase preference; GpT and TpG dinucleotides occur more frequently and are often tandemly arranged. Empirically, the data do not allow for prediction of which strand may be poised to be methylated. The results with the mammalian DCMase, which suggest sequence-dependent binding affects for a 26
5 base-pair expanse (or more), are quite reasonable given the DNA footprinting results for the bacterial enzymes mentioned. The binding asymmetry suggested by the results herein was likely induced by the design of the starting population, because one strand was guanine-rich while the other was cytosine-rich. This design was chosen in order to avoid introducing multiple CpG dinucleotides that could complicate the assessment
10 of flanking sequence contributions around a single CpG.

DCMase interactions with DNA are influenced by helical geometries

Dinucleotide analysis has been useful for understanding sequence-dependent conformational parameters of DNA (El Hassan, M.A. and Calladine, C.R., 1996,
15 Propeller-twisting of base-pairs and the conformational mobility of dinucleotide steps in DNA, J. Mol. Biol. 259:95-103; Hunter, C.A., 1993, Sequence-dependent DNA structure. The role of base stacking interactions, J. Mol. Biol. 230:1025-1054; Yanagi, K., et al., 1991, Analysis of local helix geometry in three B-DNA decamers and eight dodecamers, J. Mol. Bio. 217:201-214). The crystallography-derived
20 parameters are generally similar for the protein bound and free states (Calladine, C.R. and Drew, H.R., 1996, A useful role for "static" models in elucidating the behaviour of DNA in solution, J. Mol. Biol. 257:479-485). Dinucleotide conformational parameters have a limited range which are dependent on the two nucleotides immediately flanking the dinucleotide step in question. Because there are 136 four-
25 base steps, the understanding of the sequence-dependent helix geometry at this level is still incomplete (El Hassan & Calladine, 1996, *supra*; Yanagi et al., 1991, *supra*). More distant nucleotides also have significant effects on CpG helical parameters (Lefebvre, A., et al., 1996, Solution structure of the CpG containing d(CTTCGAAG)₂ oligonucleotide: NMR data and energy calculations are compatible with a BI/BII
30 equilibrium at CpG. Biochemistry 35:12560-12569).

These analyses provide the basis for a qualitative interpretation of DNA conformational features important for the stabilization of the initial DCMase:DNA complex. Guanosine-rich stretches, best represented in this Example by the GC-box
35 and the selected 5'-regions, often assume an A-DNA conformation. Guanosine-rich helices are under-wound because neighboring guanine bases tend to overlap and lead

to low dinucleotide twist angles (McCall, M., et al., 1985, The crystal structure of d(G-G-G-G-C-C-C-C). A model for poly(dG):poly(dC). J. Mol. Biol. 183:385-396; Yanagi et al., 1991, *supra*; El Hassan & Calladine, 1996, *supra*). Also, base-pair slide is allowed more freely in GpG than other steps, due mainly to a low propeller-twist parameter. A-DNA thus differs from B-DNA in that the minor groove is wide and shallow while the major groove is narrower and deeper. While little is known about the DCMTase:DNA interface, the enzyme contains the peptide motif ⁷¹⁶SPKK⁷¹⁹, which is found in proteins known to interact with the minor groove of DNA (Churchill, M. & Suzuki, M., 1989, "SPKK" motifs prefer to bind to DNA at A/T-rich sites, EMBO J. 8:4189-4195). The preference for sequences which have A-DNA like features may be due to DCMTase:DNA interactions mediated by this motif at some distance from the protein elements involved in CpG recognition. The GpT and TpG dinucleotide repeats, observed more frequently in the DCMTase selected 3'-flank, have unique sets of conformational parameters that can increase helical flexibility (Nagaich, A.K., et al., 1994, CA/TG sequence at the 5' end of oligo(A)-tracts strongly modulates DNA curvature, J. Biol. Chem. 269:7824-7833; Beutel, B.A. & Gold, L., 1992, *In vitro* evolution of intrinsically bent DNA, J. Mol. Biol. 228:803-812; Lyubchenko, Y.L., et al., 1993, CA runs increase DNA flexibility in the complex of Cro Protein with the O_R3 site, Biochemistry 32:4121-4127; Haniford, D.B. & Pulleybank, D.E., 1983, Facile transition of poly[d(TG):d(CA)] into a left-handed helix in physiological conditions, Nature 302:632-634).

Like the TpG step, CpG is considered "malleable" because the local conformations are dependent on flanking base-pairs (Lefebvre, A., et al., 1996, Biochemistry 35:12560-12569; Lefebvre, A. Mauffet, O., Hartmann, B., Lescot, E. & Fermandjian, S., 1995, Biochemistry 34:12019-12028; Hunter, C.A., 1993, J. Mol. Biol. 230:1025-1054; Prive, G.G., et al., 1991, J. Mol. Biol. 217:177-199; Grzeskowiak, K., et al., 1991, J. Biol. Chem. 266:8861-8883). Severe effects on the geometrical parameters associated with a centrally located CpG have been measured for at least 15 different sequences. The structures of two oligonucleotides containing the consensus CRE element, TGACCGTCA, have been determined (Mauffet, O., et al., 1992, J. Mol. Biol. 227:852-875; Konig, P. & Richmond, T.J., 1993, J. Mol. Biol. 233:139-154). Several sequences closely related to the GC-box consensus, GGGGCGGGGC (SEQ ID NO:3), have also been crystallized.

A small twist angle is characteristic of CpG embedded in guanine/cytosine-rich sequences and likely adds to the overall A-DNA character (Haran, T.E., et al., 1987, The crystal structure of d(CCCCGGGG): A new A-form variant with an extended backbone conformation, J. Biomol. Struct. Dynam. 5:199-217; Heinemann, U., et al., 1987, Crystal structure analysis of an A-DNA fragment at 1.8Å resolution: d(GCCCGGGC), Nucl. Acids Res. 15:9531-9549; Rabinovich, D., et al., 1988, Structures of the mismatched duplex d(G-G-G-T-G-C-C-C) and one of its Watson-Crick analogues d(G-G-G-C-G-C-C-C), J. Mol. Biol. 200:151-161; Verdaguer, N., et al., 1991, Molecular structure of a complete turn of A-DNA, J. Mol. Biol. 221:623-635; Frederick, C.A., et al., 1989, Molecular structure of an A-DNA decamer d(ACCGGCCGGT), Eur. J. Biochem. 181:295-307; Conner, B.N., et al., 1984, Helix geometry and hydration in an A-DNA tetramer: CCGG. J. Mol. Biol. 174:663-695; McCall et al., 1985, The crystal structure of d(G-G-G-G-C-C-C-C). A model for poly(dG):poly(dC), J. Mol. Biol. 183:385-396). Conversely, adenine/thymine-rich flanking sequences can lead to negative roll and high twist values at the CpG, so that the helix conforms more to B-DNA (Lefebvre, A., et al., 1996, Solution structure of the CpG containing d(CTTCGAAG)₂ oligonucleotide: NMR data and energy calculations are compatible with a BI/BII equilibrium at CpG, Biochemistry 35:12560-12569; Mauffet, O., et al., 1992, The fine structure of two dodecamers containing the cAMP responsive element sequence and its inverse, J. Mol. Biol. 227:852-875; Grzeskowiak, K., et al., 1991, The structure of B-helical CGATCGATCG (SEQ ID NO:6) and comparison with CCAACGTTGG (SEQ ID NO:4), J. Biol. Chem. 266:8861-8883; Prive, G.G., et al., 1991, Structure of the B-DNA decamer C-C-A-A-C-G-T-T-G-G (SEQ ID NO:7) and comparison with isomorphous decamers C-C-A-A-G-A-T-T-G-G (SEQ ID NO:5) and C-C-A-G-G-C-T-G-G, J. Mol. Biol. 217:177-199; Bingman, C.A., et al., 1992, Crystal and molecular structure of the A-DNA dodecamer d(CCGTACGTACGG (SEQ ID NO:8)), J. Mol. Biol. 227:738-756). The backbone torsion angles that connect the cytidine and guanosine residues in these structures are particularly interesting. The large slide associated with extensive inter-strand guanine stacking tends to stretch and contort the a and g torsion angles into the B_{II} conformation (Haran, T.E., et al., 1987, The crystal structure of d(CCCCGGGG): A new A-form variant with an extended backbone conformation, J. Biomol. Struct. Dynam. 5:199-217; Rabinovich et al., 1988, *supra*; Lefebvre et al., 1996, *supra*; El Antri, S., et al., 1993, Structural deviations at CpG provide a plausible explanation for the high frequency of mutation at this site, J. Mol. Biol. 230:373-378). B_{II} has an unusual *trans*, *trans* arrangement of

a and g torsion angles that is most often associated with A-DNA. B_{II} may be more readily attained by a CpG with guanine/cytosine-rich flanking sequences than with adenine/thymine-rich ones. Mechanically speaking, the B_{II} conformation allows for a crankshaft motion to modulate a destacking of bases (Haran et al., 1987, *supra*). This is likely an early event in the base flipping process mediated by DNA methyltransferases (Allan, B.A. & Reich, N.O., 1996, Targeted base stacking disruption by the EcoRI DNA methyltransferase, *Biochemistry* 35:14757-62).

The functional importance of the CpG phosphate orientation and flexibility, and DCMTase:phosphate interactions in general, have been studied using the *M.HhaI*:DNA cocrystal structure (Klimasauskas, S., et al., 1994, *HhaI* methyltransferase flips its target base out of the DNA helix, *Cell* 76:357-369; Cheng X; Blumenthal RM., 1996, Finding a basis for flipping bases, *Structure* 4:639-645). This structure has the target cytosine positioned outside of the helical cylinder covalently trapped by the enzyme. Surprisingly few contacts are made directly with the bases and extensive interactions with the backbone are asymmetrically located around the extrahelical cytosine. Interactions with the two phosphates on the 5'-side of this cytosine appear to be particularly important (5'-²pG³pC⁴pG⁵pC⁶p-3') and only phosphates 2 through 5 show several angstrom displacement when compared to the uncomplexed DNA. The peptide regions which contact the phosphates are conserved among numerous bacterial cytosine DNA methyltransferases (Cheng & Blumenthal, 1996, *supra*). For *M.HhaI*, phosphate ³p is contacted by Arg¹⁶⁵ and Ser⁸⁵, and sequence alignment suggests that Arg¹³¹⁵ and Ser¹²³³ may play analogous roles in the mammalian DCMTase. Also, Arg⁹⁸ which contacts ⁵p and Lys⁹⁰ which contacts ⁶p in *M.HhaI* have homologous residues in the DCMTase, namely Lys¹²⁴⁵ and Arg¹²³⁷.

The murine DCMTase has a DNA binding specificity that is similar to the catalytic specificity. The preference of the enzyme for guanine/cytosine-rich sequences may reflect a preferred positioning of backbone phosphates within the DCMTase:DNA complex. DCMTase may use the specificity advantage in localizing to certain genomic regions or to preferentially methylate guanine/cytosine-rich DNA *in vivo*. The function of methylation in bacteria as a primitive immune system, may be a major function for the eukaryotic methyltransferases. Many human viruses are very guanine/cytosine-rich and the discrimination we identified may aid in the specific deactivation of infected viral DNA.

Example 2: Kinetic Mechanism and Identification of a Potent Inhibitor of Murine DNA Cytosine-C⁵ Methyltransferase

This example provides four types of steady-state kinetic analyses to identify the order of substrate addition to the enzyme and the order of product release. In addition, this
 5 example identifies a potent single-stranded DNA inhibitor of DCMTase.

Materials

S-adenosyl-L-[methyl-³H]methionine (75 Ci/mmol, 1 mCi/ml, 1 Ci=37 GBq) was from Amersham Corporation. Unlabeled AdoMet (Sigma Chemical Company, St.
 10 Louis, MO) was further purified as described (Flynn, J., et al., 1996, Biochemistry 35:7308-7315). Routinely, 125 mM AdoMet stocks were prepared at a specific activity of 5.8×10^3 cpm/pmol. Two lots of poly(dI-dC:dI-dC) were purchased from Pharmacia Biotech, Inc. (Piscataway, NJ) with an average length of 6250 and 5000 base pairs. DE81 filters were purchased from Whatman, Inc. Other standard chemicals
 15 and reagents were purchased from Sigma Chemical Company or Fisher Scientific (Hampton, New Hampshire).

DNA cytosine C-5 methyltransferase was purified from mouse erythroleukemia cells as described (Xu, G., et al., 1995, Purification and Stabilization of mouse DNA
 20 methyltransferase, Biochemi.Biophys.Rev.Communi 207:544-551). Two separate preparations, with concentrations of 380 nM and 260 nM, were confirmed to have equivalent activities with the substrates studied.

DNA Substrate Preparation

25 The following three oligonucleotides mimic the GC-box transcriptional *cis*-regulatory element, in bold type, and were prepared as described (Flynn et al., 1996, *supra*). The central CpG is underlined.

GC-box a: 5'-GGGAATTCAAGGGG**CG**GGGCAAGGATCCAG-3'
 30 (SEQ ID NO:9)
 GC-box b: 5'-CTGGATCCTTGCCCC**CG**CCCCCTTGAATTCCC-3' (SEQ ID NO:10)
 GC-box b^{MET}: 5'-CTGGATCCTTGCCCC**mCG**CCCCCTTGAATTCCC-3'

Steady-State Kinetic Assays

Duplicate 25 μ L reaction volumes contained 3.0 nM DCMTase and 10 μ M AdoMet in MR buffer (100 mM Tris-HCl, pH 8.0, 10 mM EDTA, 200 μ g/ml BSA, 10 mM DTT). After preincubation at room temperature for up to 10 minutes, reactions were initiated by the addition of poly(dI-dC:dI-dC) and, if indicated, inhibitor DNA or reaction products. In several experiments it was found that initiating a reaction containing DNA with AdoMet yielded similar results to those routinely used. Single-stranded DNA was heated to 90 °C and quick cooled on ice, prior to initiation of the reaction. Freeze-thawed DNA produced equivalent results. Incubations were at 37 °C for 60 minutes. The poly(dI-dC:dI-dC) concentrations were 2.0, 4.0, 8.0, 16, 35, 80, 160, 250, 400, 700 and 1000 pM. In some experiments, GC-box a/b was the substrate, using 100 nM DCMTase and DNA concentrations of 0.20, 0.40, 1.0, 2.0, 4.0, 8.0, 15, 23 and 35 μ M. The reaction was stopped after 60 minutes by transferring 20 μ L of the reaction onto a DE 81 filter paper that was processed as described (Flynn et al., 1996). The radioactivity above the background, determined from assays without added poly(dI-dC:dI-dC), was converted to initial velocities and expressed as picomoles of methyl groups transferred to poly(dI-dC:dI-dC) per hour and plotted in double reciprocal form. The substrates poly(dI-dC:dI-dC) and AdoMet, competitor DNA and reaction product concentrations were varied as indicated in the Figures.

For double reciprocal plots of velocity versus substrate concentration (Figures 12A & 12B), reactions contained 20 nM DCMTase in 100 mM Tris pH 8.0, 10 mM EDTA, 10 mM DTT, 200 μ g/mL BSA. Incubations were at 37 °C for 60 minutes. For Figure 12A, poly(dI-dC:dI-dC) was varied at: triangles, 4 μ M; squares, 2 μ M; diamonds, 1 μ M; circles, 0.5 μ M, while AdoMet was constant. For Figure 12B, AdoMet was varied at: triangles, 112 pM; squares, 56 pM; diamonds, 28 pM; circles, 14 pM, while poly(dI-dC:dI-dC) concentration remained constant.

For the double reciprocal plot of velocity versus poly(dI-dC:dI-dC) with varying GC-box b concentrations (Figure 13), reactions contained 3.0 nM DCMTase and 10 μ M AdoMet in 100 mM Tris pH 8.0, 10 mM EDTA, 10 mM DTT, 200 μ g/mL BSA. The poly(dI-dC:dI-dC) concentrations were 10, 13, 20, 40 and 100 pM. The GC-box b concentrations were: diamonds, 0; circles, 0.75 μ M; triangles, 1.5 μ M; squares, 5.0 μ M. Incubations were at 37 °C for 60 minutes.

-41-

For the double reciprocal plot of velocity vs. poly(dI-dC:dI-dC) with varying GC-box b^{MET} concentrations (Figure 14), reactions contained 2.0 nM DCMTase and 10 μ M AdoMet in 100 mM Tris pH 8.0, 10 mM EDTA, 10 mM DTT, 200 μ g/mL BSA. The poly(dI-dC:dI-dC) concentrations were 1.5, 3.0, 7.5, 15 and 20 pM. The GC-box b^{MET} concentrations were: squares, 0; circles, 10 nM; diamonds, 20 nM; triangles, 40 nM. Incubations were at 37 °C for 60 minutes.

For the double reciprocal plot of velocity versus AdoMet with varying GC-box b^{MET} concentrations, (Figure 15), reactions contained 4.0 nM DCMTase and 50 pM poly(dI-dC:dI-dC) in 100 mM Tris pH 8.0, 10 mM EDTA, 10 mM DTT, 200 μ g/mL BSA. The AdoMet concentrations were 0.75, 1.5, 3.0 and 6.0 μ M. The GC-box b^{MET} concentrations were: squares, 0; circles, 20 nM; diamonds, 40 nM; triangles, 80 nM. Incubations were at 37 °C for 60 minutes.

For the double reciprocal plot of AdoHcy product inhibition with varying AdoMet concentrations (Figure 16), reactions contained 20 nM DCMTase and 40 pM poly(dI-dC:dI-dC) in 100 mM Tris pH 8.0, 10 mM EDTA, 10 mM DTT, 200 μ g/mL BSA. The AdoMet concentrations were 0.50, 1.0, 2.0, 4.0 and 8.0 μ M. The AdoHcy concentrations were: squares, 0; diamonds, 0.75 μ M; circles, 1.5 μ M; triangles, 3.0 μ M; notched squares, 6.0 μ M. Incubations were at 37 °C for 60 minutes.

For the double reciprocal plot of AdoHcy product inhibition with varying poly(dI-dC:dI-dC) concentrations (Figures 17A-C), reactions contained 20 nM DCMTase in 100 mM Tris pH 8.0, 10 mM EDTA, 10 mM DTT, 200 μ g/mL BSA. Incubations were at 37 °C for 60 minutes. The poly(dI-dC:dI-dC) concentrations were 2.5, 5.0, 10, and 20 pM. The AdoHcy concentrations were: squares, 0; diamonds, 15 μ M; circles, 30 μ M. For Figure 17A, AdoMet was held constant at 1.2 μ M. For Figure 17B, AdoMet was held constant at 8 μ M. Figure 17C shows secondary slope replots from another series of experiments in which the AdoMet concentrations were: circles, 6.3 μ M; diamonds, 2.5 μ M; squares 1 μ M.

For the double reciprocal plot of poly(dId^mC:dId^mC) product inhibition with varying AdoMet concentrations (Figure 18), reactions contained 20 nM DCMTase and 60 pM poly(dI-dC:dI-dC) in 100 mM Tris pH 8.0, 10 mM EDTA, 10 mM DTT, 200 μ g/mL BSA. The AdoMet concentrations were 1.0, 2.0, 4.0 and 8.0 μ M. The poly(dId^mC:dId^mC) concentrations were: squares, 0; diamonds, 5.0 pM; circles, 10

pM; triangles, 20 pM. Incubations were at 37 °C for 60 minutes. Experimental data are shown scattered around lines derived from a fit to equation 5 for noncompetitive inhibition.

- 5 For the double reciprocal plot of poly(dId^mC:dId^mC) product inhibition with varying poly(dI-dC:dI-dC) concentrations (Figures 19A-B), reactions contained 20 nM DCMTase and 1.5 μM AdoMet in 100 mM Tris pH 8.0, 10 mM EDTA, 10 mM DTT, 200 μg/mL BSA. Incubations were at 37 °C for 60 minutes. The poly(dI-dC:dI-dC) concentrations were 20, 40, 80, 120 and 160 pM. The poly(dId^mC:dId^mC)
- 10 concentrations were: squares, 0; triangles, 34 pM; circles, 45 pM; diamonds, 68, notched squares, 90 pM. Experimental data are shown scattered around lines. In Figure 19A, lines are derived from a fit to equation 4 for competitive inhibition. In Figure 19B, lines are derived from a fit to equation 5 for noncompetitive inhibition. Both fittings are not acceptable.
- 15 For initial velocity plots of different poly(dIdC:dIdC) lengths (Figure 20), reactions contained 20 nM DCMTase and 10 μM AdoMet in 100 mM Tris pH 8.0, 10 mM EDTA, 10 mM DTT, 200 μg/mL BSA. Incubations were at 37 °C for 60 minutes. The poly(dI-dC:dI-dC) sizes were: circles, 100 base-pairs; diamonds, 500 base-pairs;
- 20 triangles, 2000 base-pairs; squares, 5000 base-pairs. The inset provides a zoom in along the x-axis toward the origin to show the quality of the data.

Fragmentation of poly(dI-dC:dI-dC)

- 25 Sonication was used to break a 5000 base-pair average length poly(dI-dC:dI-dC) to lengths of approximately 2000, 1400, 600, 500 and 100 base-pairs using a Branson Sonifier 450 with a microbore tip. Lengths were estimated by agarose gel electrophoresis using DNA size standards.

Preparation of poly(dId^mC:dId^mC)

- 30 Poly(dI-dC:dI-dC) was methylated to completion with M.SssI (New England Biolabs). The methylation reaction was optimized and the apparent K_m^{DNA} was determined to be 0.40 nM for M.SssI using 6250 base-pair poly(dI-dC:dI-dC). For reaction efficiency and sufficient yields, a 500 μL reaction contained 1.0 nM poly(dI-dC:dI-dC). AdoMet was added to 100 μM to provide an excess level of methyl-groups to complete the
- 35 reaction. Three 20 unit aliquots of M.SssI were added every 10 hours in MR buffer.

After cleaning the DNA by standard organic extraction methods, it was resuspended to 10 nM in TE (10mM Tris pH 8.0, 1mM EDTA) and subjected to the methylation reaction using *M.SssI* and radiolabeled AdoMet. Background, 230 cpm, was detected with this preparation at 0.80 nM and a similar control experiment with 0.80 nM
5 unmethylated poly(dI-dC:dI-dC) generated 37,000 cpm. The methylated DNA, poly(dI-d^mC:dI-d^mC), was resistant to digestion by *HhaI* endonuclease and the control DNA was digested to small fragments, as determined by agarose gel electrophoresis.

Isotope Partitioning Analysis

10 A pre-steady-state approach was used to determine the catalytic competency of the DCMTase:AdoMet complex. The complex was formed at 37 °C using 20 nM DCMTase and tritiated AdoMet at a concentration of 10 µM. The reaction was initiated by adding a mixture of 400 pM poly(dI-dC:dI-dC) and 100 mM unlabeled AdoMet. After a one hour incubation at 37 °C, the reactions were treated as stated
15 above.

Molecular Partitioning Analysis

A pre-steady-state approach was used to determine the catalytic competency of the DCMTase:DNA complex. Two different sizes of substrate DNA, 1400 and 600 base-
20 pair poly(dI-dC:dI-dC), were used to distinguish if the initial complex proceeded in the forward direction or dissociated before DCMTase performed chemistry. The complex was formed at 37 °C for 1.5 minutes with 5 nM DCMTase and the 1400 base-pair poly(dI-dC:dI-dC) at 0.20 nM, then a mixture containing 2.0 µM tritiated AdoMet (neat stock concentration, 13 µM) plus an excess of the molecular competitor, 600
25 base-pair poly(dI-dC:dI-dC) at 5.0 nM was added to initiate catalysis. Aliquots were removed at 1.5, 3 and 9 minutes followed by centrifugation through a P-6 spin column (Bio-Rad) to trap unincorporated label. DNA were separated on an 6% polyacrylamide, 8M urea gel run at 400 V for 4.5 hours. Standard methods of fluorography were used with LiquiScint (National Diagnostics) as the fluor. The dried
30 gel was exposed to Fuji XAR film for three months at -70 C.

Data Analysis

The Michaelis-Menton equation was used for studies into DNA length contributions to catalysis using KaliedaGraph 2.1.2 (Synergy Software). For mechanistic

determinations, the nomenclature used is that of Cleland, W.W., 1963a. *Biochim. Biophys. Acta* 67:104–137. All steady-state data were analyzed using regression analyses of the appropriate initial velocity equation, listed below, using the Cleland programs (Cleland, W.W., 1979, *Statistical analysis of enzyme kinetic data*, Methods in Enzymol. 63:103–138).

Substrate Inhibition:

$$v = \frac{VA}{K_a + A + A^2/K_i} \quad (1)$$

Sequential Mechanism:

$$v = \frac{VAB}{K_{ia}K_b + K_aB + K_bA + AB} \quad (2)$$

Ping Pong Mechanism:

$$v = \frac{VAB}{K_aB + K_bA + AB} \quad (3)$$

Competitive Inhibition:

$$v = \frac{VA}{K_a(1 + K_{is}) + A} \quad (4)$$

Noncompetitive Inhibition:

$$v = \frac{VA}{K_a(1 + K_{is}) + A(1 + K_{ii})} \quad (5)$$

Uncompetitive Inhibition:

$$v = \frac{VA}{K_a + A(1 + K_{ii})} \quad (6)$$

The algorithms perform a non-linear least squares fit to the entire data set. Mechanistic determinations were made by comparison of the sigma values associated with the fit to each equation. The standard errors associated with fitted parameters and graphical analysis of the experimental data scattered around the calculated best fit lines were also considered in making an assignment.

Preparation of poly(dI-d^mC:dI-d^mC)

Poly(dI-dC:dI-dC) was methylated to completion with M.SssI (New England Biolabs, Beverly, Massachusetts). Optimization of the methylation reaction was investigated and the apparent K_m^{1C} was determined to be 0.4 nM for M.SssI. For reaction efficiency and sufficient yields, a 500 mL reaction contained 1 nM poly(dI-dC:dI-dC). AdoMet was added to 100 μ M to provide an excess level of methyl-groups to complete the reaction. Three 20 unit additions of M.SssI were done every 10 hours in our methylation buffer. Complete methylation of poly(dI-dC:dI-dC) was tested. After cleaning the DNA by standard methods, it was resuspended to 10 nM in TE and subjected to the methylation reaction using M.SssI and radiolabeled AdoMet. Background, 230 cpm, was detected with this preparation at 0.8 nM and a control experiment generated 37,000 cpm. The methylated DNA, poly(dI-d^mC:dI-d^mC), was resistant to digestion by HhaI endonuclease and the control DNA was digested to small fragments, determined by agarose gel electrophoresis.

Fragmentation of poly(dIdC:dIdC)

Sonication was used to break a 5000 base-pair average length poly(dIdC:dIdC) to sizes of approximately 2000, 500 and 100 base-pairs. The sizes were determined by agarose gel electrophoresis and comparison to DNA size standards.

Results

The DNA substrate used in the steady-state studies was poly(dI-dC:dI-dC). This substrate contains tandem methylation sites in which guanosine has been replaced by inosine. Methylation is catalyzed at a higher rate with this substrate than with other DNA (Flynn, J., et al. (1996) Biochemistry 35:7308-7315; Pedrali-Noy, G., & Weissbach, A., (1986) J. Biol. Chem. 261:1, 7600-77602). Poly(dI-dC:dI-dC) provides a uniform sequence and limits the potential complexities found with large

cloned sequences that contain many randomly situated CpG dinucleotides, each having different flanking sequence contributions to binding and catalysis (Flynn et al., 1996 *supra*).

5 *Substrate Inhibition*

Linn and coworkers reported that high concentrations of large DNA resulted in DCMTase inhibition, and proposed that multimeric forms of the enzyme are required for activity (Hitt, M.M., et al., 1988, J. Biol. Chem. 263:4392-4399). To distinguish between this and alternative explanations, a short 30 base-pair DNA substrate was
 10 tested for inhibition at high concentrations. Figure 11 shows the initial velocity results for poly(dI-dC:dI-dC), 6250 base-pairs, and GC-box a/b in terms of reduced concentrations (S/K_m^{DNA}). Initial velocity data for both substrates were fit to equation 1, which is a standard equation for analyzing substrate inhibition. K_m^{DNA} was determined to be 5.5 ± 0.9 pM and 0.31 ± 0.13 μ M, and K_i^{DNA} was 1010 ± 170
 15 pM and 43 ± 22 μ M for poly(dI-dC:dI-dC) and GC-box a/b, respectively (Table 3). In both cases, DNA concentrations greater than 20 times K_m^{DNA} caused substrate inhibition. AdoMet utilization was calculated to be less than 0.5%. Product inhibition by S-adenosyl homocysteine formation is therefore unlikely. AdoMet substrate inhibition was not observed at concentrations up to 30 times K_m^{AdoMet} . The substrate
 20 inhibition observed implicates a second DNA molecule binding to the enzyme and inhibiting catalysis.

Table 3: Substrate Inhibition Constants^a

25	GC-box a/b	43000 \pm 22000 nM
	Poly(dIdC:dIdC)	1.0 \pm 0.2 nM

^a The constants reported, K_i in equation 1, were derived from non-linear regression to the appropriate rate equations as described above. The nomenclature is that of
 30 Cleland (1963b).

Initial Velocity Studies with Poly(dI-dC:dI-dC) and AdoMet

Double reciprocal plots of initial velocity versus the substrate concentrations are shown in Figure 12. DNA concentrations near K_m were used to avoid non-Michaelis
 35 behavior (see substrate inhibition studies above). The transformed data were best fit by lines intersecting left of the y-axis using a non-linear regression of equation 2, a

standard equation for analyzing the steady-state mechanism. The true Michaelis constants derived were $K_m^{pdIdC} = 36 \pm 5$ pM and $K_m^{AdoMet} = 1.4 \pm 0.2$ μ M. The intersecting patterns rule out a nonsequential mechanism and implicate a sequential order of substrate addition in which both DNA and AdoMet add to the enzyme surface before products are released. However, a prudent assignment of a kinetic mechanism requires additional kinetic arguments.

Dead-End Inhibition with Single-Stranded DNA

A previous kinetic study showed no detectable enzyme activity with single-stranded GC-box a and GC-box b (Flynn, J., et al., 1996, Murine DNA cytosine-C5 methyltransferase: Pre-steady- and steady-state kinetic analyses with regulatory DNA sequences, *Biochemistry* 35:7308-7315). In contrast, the DCMTase binds these same oligonucleotide substrates with affinities comparable to those of other DNA (see Example 1). For these reasons it was presumed that single-stranded GC-box substrates could act as dead-end inhibitors of the reaction with poly(dI-dC:dI-dC). Dead-end inhibitors can provide a strong methodology for assessing whether a kinetic mechanism is random or ordered. Inhibition of poly(dI-dC:dI-dC) methylation by single-stranded GC-box b was studied at 15 μ M AdoMet. The data were best fit by equation 5, a standard equation for noncompetitive inhibition, and generated the intersecting double reciprocal pattern shown in Figure 13. The inhibition constants were determined to be $K_{is} = 3.6 \pm 1.5$ μ M and $K_{ij} = 6.8 \pm 1.2$ μ M (Table 4). K_{ij} is the inhibition constant associated with an intercept effect and K_{is} is the inhibition constant associated with a slope effect from families of double reciprocal plots. An alpha factor, K_{ij}/K_{is} , of 1.9 was determined and suggests that the partitioning of this inhibitor slightly favors addition to the free enzyme over the DCMTase:poly(dI-dC:dI-dC) intermediate.

Table 4: Dead-End Inhibition Constants and Mode of Inhibition^a

	DNA	Inhibition Constant (nM)	Mode of Inhibition
	GC-box b	3600 +/- 1500 ^b	Noncompetitive with poly (dIdC:dIdC)
5	GC-box b	6800 +/- 1200 ^c	Noncompetitive with poly (dIdC:dIdC)
	GC-box b ^{MET}	20 +/- 3 ^d	Uncompetitive with poly (dIdC:dIdC)
	GC-box b ^{MET}	25 +/- 10 ^e	Competitive with Adomet

^aThe constants reported were derived from non-linear regression to the appropriate rate equations as described above. The nomenclature is that of Cleland (1963b).

^bThe inhibition constant refers to the slope derived K_{ij} in equation 5.

^cThe inhibition constant refers to the intercept derived K_{ij} in equation 5.

^dThe inhibition constant refers to the intercept derived K_{ij} in equation 6.

^eThe inhibition constant refers to the slope derived K_{ij} in equation 4.

15 The CpG methylated homolog of GC-box b, GC-box b^{MET}, was studied for inhibition under similar conditions. The data were best fit by the log form of equation 6, a standard equation for uncompetitive inhibition. The inhibition constant, K_{ij} , was estimated to be 20 +/- 3 nM. The double reciprocal transformation is shown in Figure 20 14. Remarkably, a single 5-^mC substitution appears responsible for a 200-fold lower inhibition constant and a change in the mode of inhibition. The uncompetitive nature of inhibition suggests that GC-box b^{MET} and poly(dI-dC:dI-dC) bind to distinct sites on the DCMTase surface and that poly(dI-dC:dI-dC) binds prior to GC-box b^{MET}.

25 Another characterization of the potent inhibition observed with GC-box b^{MET} was obtained by varying AdoMet and GC-box b^{MET} concentrations using a constant 50 pM poly(dI-dC:dI-dC). The data from initial velocities were best fit to equation 4, which is a standard equation for competitive inhibition. The estimated inhibition constant was $K_{is} = 25 +/- 10$ nM. The intersection of the fit lines on the 1/velocity 30 axis in Figure 15 suggests that GC-box b^{MET} and AdoMet bind competitively to the same poly(dI-dC:dI-dC)-bound form of the enzyme.

The two inhibition constants determined for GC-box b^{MET} are in good agreement at about 20 nM. The patterns observed provide strong evidence for an ordered Bi-Bi 35 kinetic mechanism with substrate DNA binding to the enzyme first and AdoMet binding second, followed by the release of products (Spector & Cleland, 1981,

Meanings of K_i for conventional and alternative-substrate inhibitors, Bio. Pharm. 30:1-7). In the absence of poly(dI-dC:dI-dC), GC-box b^{MET} bound free enzyme with a 120-fold lower affinity (see Example 1).

5 *Product Inhibition Studies*

Product inhibition studies were pursued to further identify the steady-state kinetic mechanism (Table 5). The DCMTase reaction product AdoHcy was a competitive inhibitor of AdoMet. The competitive nature of AdoHcy with respect to AdoMet binding, $K_{is} = 1.4 \pm 0.2 \mu M$, suggests that AdoMet and AdoHcy bind to the same
10 form of the enzyme (Figure 16) or that the kinetic mechanism is Theorell-Chance. The Theorell-Chance kinetic mechanism is a simplification of the Ordered B-Bi scheme.

Table 5: Product Inhibition of Murine DNA Cytosine-C⁵ Methyltransferase.^a

15

Product	Varied Substrate	Fixed Substrate	Type of Inhibition	Inhibition Constant ^b
¹ mC	IC	AdoMet	NC	nd ^c
20 ¹ mC	AdoMet	IC	NC	K _{is} 5.3 \pm 2.1 pM K _{ii} 30 \pm 12 pM
AdoHcy	IC	AdoMet	NC/UC	Nd ^d
AdoHcy	AdoMet	IC	C	K _{is} 1.4 \pm 0.2 μM


25 ^a ¹mC, fully methylated poly(dIdC:dIdC); IC, poly(dIdC:dIdC); AdoHcy, S-adenosyl homocysteine; AdoMet, S-adenosyl methionine; C, competitive; NC, noncompetitive; UC, uncompetitive inhibition.

^b K_{is} refers to the inhibition constant derived from a slope affect. K_{ii} refers to the inhibition constant derived from an intercept affect.

30 ^c nd, not determined. The determination of an inhibition constant may be complicated by binding to a second nucleic acid binding site on the DCMTase.

^d nd, not determined. The inhibition constants are dependent on the fixed AdoMet concentration and inhibition is not overcome by saturating AdoMet concentrations. Instead, the inhibition profiles are noncompetitive-like at low AdoMet and
35 uncompetitive-like at high AdoMet concentrations.

A distinctive inhibition profile is revealed with varying AdoHcy and poly(dIdC:dIdC). Two families of plots were obtained with AdoMet at different constant levels, 1.2 μ M and 8 μ M (Figures 17A and 17B, respectively). Increasing the AdoMet concentration had the gradual effect of changing the plots from a noncompetitive-like pattern at concentrations near K_m^{AdoMet} to an uncompetitive pattern at higher AdoMet concentrations. There is a decrease in scale of the y-axis in Figure 17B compared to 17A and a lack of a significant change of the data collected at high poly(dIdC:dIdC) concentrations, the points closest to the y-axis. The data at low AdoMet concentrations fitted slightly less well to an uncompetitive model, $K_{ii} = 2.0 \pm 0.6$ μ M, than to a noncompetitive model that produced the constants $K_{is} = 63 \pm 71$ μ M and $K_{ii} = 2.5 \pm 1.0$ μ M. The K_i^{AdoHcy} was independently determined in Figure 15 to be 1.4 μ M. The slope and y-intercept replots from each AdoHcy versus poly(dIdC:dIdC) series were all linear. Another study confirmed these results and showed a gradual effect of going from a noncompetitive to an uncompetitive model using three AdoMet concentrations; 1, 2.5, and 6.3 μ M. This analysis demonstrates that the slope contribution to AdoHcy inhibition is minimal at low AdoMet concentrations, inhibition cannot be overcome by high AdoMet concentrations, and AdoHcy binds to a different enzyme form than poly(dIdC:dIdC). This is strong evidence for an Ordered Bi Bi mechanism in which initial DNA binding is followed by AdoMet binding and that the following reaction step is irreversible.



Also, the last product to leave the enzyme cannot be AdoHcy if poly(dIdC:dIdC) is the first substrate to bind DCMTase. Uncompetitive inhibition with AdoHcy and DNA was also observed with *M.HhaI*, it provided evidence that a *M.HhaI*:DNA:AdoHcy complex can form and ruled out a catalytically significant *M.HhaI*:AdoHcy complex (Wu & Santi, 1987).

30 *Product Inhibition with Poly(dId^mC:dId^mC)*

Fully methylated poly(dId^mC:dId^mC) was prepared and used as a product inhibitor of the DCMTase reaction. Poly(dId^mC:dId^mC) was linear noncompetitive with AdoMet when poly(dIdC:dIdC) was held constant (Figure 18; Table 3). The estimated inhibition constants were $K_{is} = 5.3 \pm 2.1$ pM and $K_{ii} = 30 \pm 12$ pM. The noncompetitive pattern with AdoMet supports many different mechanisms including one in which DNA binding occurs prior to AdoMet binding.

The double reciprocal pattern for methylated DNA product versus DNA substrate would be expected to be competitive in a standard ordered Bi-Bi kinetic mechanism where DNA adds first and methylated DNA leaves last from the catalytically competent enzyme surface. The double reciprocal data obtained on five experiments appeared to be noncompetitive. However, when subjecting the data to fitting by the competitive and the noncompetitive models, graphical analysis showed that fitting in both cases was not acceptable (Figure 19). This was true for plots obtained with AdoMet concentrations held constant from 1.25 to 12.5 μM . The sensitivity of inhibition was notably abrupt, as poly(dId^mC:dId^mC) had little effect at 10 pM and completely inhibited the reaction at 100 pM. The results from one experiment are shown in Figures 23A-C with idealized lines intersecting left of the y-axis. Secondary slope and y-intercept replots (Figures 23B and 23C) were obtained and both were parabolic concave upward. This explains the difficulty in fitting the simple model and is indicative of poly(dI-d^mC:dI-d^mC) binding at two points in the catalytic cycle. Furthermore, it is evidence that a DCMTase:DNA:DNA complex can be formed. Additional steady-state kinetic experiments also support the existence of an inhibitory DCMTase:DNA:DNA complex.

20 *Isotope Partitioning Analysis with AdoMet*

Isotope partitioning analysis is a powerful strategy used to identify catalytically competent enzyme:substrate complexes (Rose, 1980; Reich & Mashhoon, 1991). The DCMTase:AdoMet complex formed with 10 μM radiolabeled AdoMet was not competent for catalysis, because a chase including 400 pM poly(dI-dC:dI-dC) and 100 μM unlabeled AdoMet produced no detectable activity. Substrate inhibition was not observed at high AdoMet concentrations. This is typical for an Ordered Bi Bi mechanism when studying the second substrate by isotope partitioning, because the DCMTase:AdoMet complex must dissociate before DCMTase can bind poly(dI-dC:dI-dC). Under these conditions, the DCMTase:poly(dI-dC:dI-dC) complex would then bind a diluted specific activity AdoMet and catalysis would not be detectable.

Molecular Partitioning Analysis

A novel assay was developed to test the competency of the initial DCMTase:poly(dI-dC:dI-dC) complex. This complex was formed with one DNA

length, 1400 base-pairs, and then challenged with an excess of smaller, 600 base-pair DNA combined with AdoMet. The initial DCMTase:poly(dI-dC:dI-dC) complex was observed to be competent for catalysis, because tritium was incorporated into the larger DNA. A control experiment allowed both DNA lengths to compete for
 5 DCMTase binding before AdoMet was added, because the smaller DNA was at a sufficiently higher concentration all of the detectable label was incorporated into it. This demonstrates that DNA, under the conditions employed, can bind first in the steady-state mechanism, and limits the assumptions made in other experiments to the Ordered Bi Bi mechanism.

10

DNA Length Contributions to Catalytic Efficiency

Sonication was used to break a 5000 base-pair average length poly(dIdC:dIdC) to sizes of approximately 2000, 500 and 100 base-pairs. Initial velocity profiles were obtained for each size (Figure 20), and the kinetic terms are compared in Table 6. A
 15 14-fold increase in K_m^{DNA} was observed as the length decreased 5000 to 100 base-pairs, but k_{cat} only dropped by one-third. The hyperbolic trend in specificity constants, k_{cat}/K_m^{DNA} (Figure 21), suggests a half maximal length of 1200 base-pairs and that lengths greater than 2000 base-pairs provide little advantage to catalytic specificity. On the contrary, DNA lengths of 500 base-pairs and smaller show a very sharp
 20 decrease in specificity. DNA length is thereby critical to maximal performance of DCMTase.

Table 6: Poly (dIdC:dIdC) length and Catalytic Efficiency^a

25	Length	Vmax (fmol hr ⁻¹)	kcat (hr ⁻¹)	Km (pM)	Kcat/Km(hr ⁻¹ M ⁻¹ x 10 ¹⁰)
	5000	12500 +/- 1100	31.2	140 +/- 30	22.2
	2000	9950 +/- 390	24.9	125 +/- 17	19.9
	500	9120 +/- 290	22.8	300 +/- 30	7.65
	100	8600 +/- 210	21.5	1890 +/- 150	1.13

30

^aConstants were determined from initial velocity analysis using the Michaelis-Menton equation.

Reciprocal plots of both substrates, AdoMet and 100 base-pair poly(dIdC:dIdC), were
 35 generated. The patterns observed were much like that shown in Figure 12 with 6250 base-pair poly(dIdC:dIdC). Although large effects in the kinetic terms were observed

with decreasing poly(dIdC:dIdC) length, the mechanism of catalysis does not appear to be affected.

Discussion

5 The data presented herein clarify some of the basic aspects of how cytosine C-5 methylation is catalyzed and perhaps controlled in eukaryotes. The order of substrate binding appears to be DNA followed by AdoMet and the order of product release appears to be AdoHcy followed by methylated DNA. Three kinetic methodologies support our assignments: initial velocity studies varying both substrates, dead-end
10 inhibition, and product inhibition. DNA substrate inhibition was common to both small, single CpG containing DNA and large, multi-site DNA. A second nucleic acid binding region on the DCMTase, distinct from the active site, is implicated from both the substrate inhibition and the dead-end inhibition studies.

15 *DCMTase Multimerization and Substrate Inhibition*

Several of the results bear directly on the previously proposed formation of reversible, multimeric complexes (Reale, A., et al., 1995, DNA binding and methyl transfer catalysed by mouse DNA methyltransferase, *Biochem. J.* 312:855-861) and the inhibition of DCMTase activity at high DNA concentrations (Hitt, M.M., et al., 1988,
20 *De novo* and maintenance DNA methylation by a mouse plastytoma cell DNA methyltransferase, *J. Biol. Chem.* 263:4392-4399). An understanding of the functional form(s) of the DCMTase is essential for future structure-function analysis, and the mechanism of DNA-mediated inhibition may be important for *in vivo* regulation of the enzyme. The DCMTase in the absence of either DNA or AdoMet
25 exists as a monomer, as determined by size exclusion chromatography (Xu, G., et al., 1995, Purification and stabilization of mouse DNA methyltransferase, *Biochemi. Biophys. Res. Communi.* 207:544-551). Active site titration suggests that the enzyme is a functional monomer (Flynn, J., et al., 1996, Murine DNA cytosine-C5 methyltransferase: Pre-steady- and steady-state kinetic analyses with regulatory
30 DNA sequences, *Biochemistry* 35:7308-7315). Further support for a 1:1 enzyme to DNA catalytic association was provided by gel mobility shift analyses (see Example 1).

- Hitt et al., 1988, *De novo* and maintenance DNA methylation by a mouse plastytoma cell DNA methyltransferase, J. Biol. Chem. 263:4392-4399) proposed that the DCMTase is inhibited at high DNA concentrations by partitioning to a monomeric enzyme bound to DNA, and that protein multimerization results in enzyme activation.
- 5 An alternative explanation could be that substrate inhibition occurs with the formation of a ternary complex (DCMTase:DNA:DNA). These models were tested with a short DNA substrate that is less likely to support protein multimerization than a long, multi-site substrate. Both substrates clearly showed inhibition at high DNA concentrations, and the normalized inhibition profiles appear very similar (Figure 11).
- 10 The corresponding substrate inhibition constants, K_i , are 150 to 180 times greater than K_m^{DNA} for these very different DNA molecules (Table 3).

The similar K_m/K_i ratios suggest that the substrate inhibition is insensitive to the number of CpG or CpI dinucleotides within the DNA. Moreover, the concentration

15 dependencies, particularly with the small DNA, show that the inhibition occurs via an intermolecular process. The results also suggest that the DCMTase has a second DNA-binding site with lower affinity for these substrates than the site involved in catalysis. The formation of an inhibitory, ternary complex that includes two DNA molecules is further supported by our inhibition studies with single-stranded DNA

20 (see below) and the existence of DNA-binding peptide motifs residing in the non-catalytic amino-terminal domain of the enzyme (Bestor, T.H., 1992, Activation of the mammalian DNA methyltransferase by cleavage of a Zn binding regulatory domain, EMBO 11:2611-2617; Chuang, L.S., et al., 1996, Characterisation of independent DNA and multiple Zn-binding domains at the N terminus of human DNA-(cytosine-

25 5_ methyltransferase: modulating the property of a DNA-binding domain by contiguous Zn-binding motifs, Chia, J., and Li, B.F.L., J. Mol. Biol. 257:935-948). Example 1 shows that DNA concentrations ten times higher than K_m produce a second less mobile band that is consistent with a second DNA binding event.

30 *Kinetic Analysis of DNA and AdoMet Binding*

Knowledge of the order of substrate binding and product dissociation is of critical importance to understanding an enzyme mechanism and the mechanisms of particular inhibitors. A first step in the kinetic characterization for DCMTase is shown in Figure

12. Several observations suggest that DNA binds first. The dead-end inhibition

35 observed in Figures 14 and 15 with GC-box b^{MET} implicates DNA (substrate) binding

prior to the inhibitor (ssDNA). These inhibition patterns are inconsistent with both a random mechanism and an ordered addition in which AdoMet must bind prior to DNA (Cleland, W.W., 1963b, The kinetics of enzyme-catalyzed reactions with two or more substrates or products II. Inhibition: Nomenclature and theory, Biochim.

5 Biophys. Acta 67:173-187). The gel shift experiments (Example 1) clearly show that DCMTase can bind DNA in the absence of AdoMet. Moreover, cofactor addition had no detectable effect on the binding affinity. While the catalytic competence of the initial binding event is uncertain, the stability of the complex is dependent on DNA sequence.

10

The product inhibition studies provided both arguments for and against the classic ordered Bi-Bi mechanism shown in Figure 22. Two studies were inconsistent with the proposed kinetic order: poly(dId^mC:dId^mC) inhibition with varied poly(dIdC:dIdC), constant AdoMet (Figure 19) and AdoHcy inhibition with varied AdoMet, constant poly(dIdC:dIdC) (Figure 16). In the first case, it is proposed that the complexities involved with a second DNA binding site have complicated the classic model in ways that are difficult to assess from just these studies. In the second case, AdoHcy exhibited competitive inhibition, but noncompetitive is expected. It may be that AdoHcy is so similar to AdoMet, in that they differ by a methyl group, that the product inhibition does not behave classically. The Theorell-Chance mechanism could also explain this result. Poly(dId^mC:dId^mC) inhibition with varied AdoMet, constant poly(dIdC:dIdC) (Figure 18) was consistent with the mechanism proposed. Also, it must be considered that the above three product inhibition studies are consistent with many different mechanisms (Segel, 1975, Enzyme kinetics behavior and analysis of rapid equilibrium and steady-state enzyme systems, John Wiley, New York, pg 653). The fourth product inhibition study, AdoHcy inhibition with varied poly(dIdC:dIdC) and constant AdoMet, appeared somewhat complicated (Figure 17A-C). Not only is the result consistent with the proposed mechanism, it is uniquely characteristic of it (Segel, 1975, *supra*).

30

An overwhelming amount of the data presented herein support a kinetic order as follows: DNA binds, then AdoMet binds and catalysis occurs, AdoHcy leaves followed by methylated DNA (Figure 22). This proposed mechanism is similar to that described by Wu & Santi (1987, Kinetic and catalytic mechanism of HhaI methyltransferase, J. Biol. Chem. 262:4778-4786) for the bacterial DCMTase, M.HhaI. We suggest that the intersecting double reciprocal plots for a rapid

35

equilibrium mechanism observed with *M.HhaI* and our observation of double reciprocal plots that intersect far from the y -axis with the murine DCMTase may be reconciled by differences in the lifetimes and partitioning of both the enzyme:DNA and enzyme:DNA:AdoMet intermediates.

5

DNA length contributes to catalytic efficiency

The investigations into poly(dIdC:dIdC) length produced some interesting findings. The apparent K_m systematically increased 14-fold when decreasing the length from 5000 to 100 base-pairs. On the contrary, Table 6 shows that k_{cat} only decreases by
10 one-third. This suggests that assembly of the competent enzyme:DNA:AdoMet complex is difficult and longer DNA promotes catalysis better than small DNA. However, once the complex is formed catalysis can proceed about as well with 100 or 5000 base-pair poly(dIdC:dIdC).

15 Facilitated diffusion of DNA binding proteins and enzymes is a well characterized phenomenon (Surby & Reich, 1996a, Contribution of facilitated diffusion and processive catalysis to enzyme efficiency: implications for the EcoRI restriction-modification system, Biochemistry 35:2201-2208, Surby & Reich 1996b, Facilitated
diffusion of the EcoRI DNA methyltransferase is described by a novel mechanism,
20 Biochemistry 35:2209-2217). It appears from these studies that DCMTase also uses facilitated diffusion to seek and stabilize the catalytic complex. The specificity constant determined of $6.1 \times 10^7 \text{ sec}^{-1} \text{ M}^{-1}$ is within an order of magnitude of the diffusion controlled limit and because this enzyme is unusually slow, k_{cat} under 30 hr^{-1} , it is expected that facilitated diffusion contributes largely to catalysis. Processivity
25 has not been addressed in our studies, however, the kinetic mechanism proposed is that expected for a processive enzyme.

Identification of a potent, reversible inhibitor

The finding that single-stranded GC-box a and GC-box b bind with reasonable
30 affinity (Example 1) was somewhat surprising given our inability to detect a significant methyl transfer activity with these sequences. The DCMTase is capable of modifying other ssDNA (Flynn, J., et al., 1996, Murine DNA cytosine-C5 methyltransferase: Pre-steady- and steady-state kinetic analyses with regulatory DNA sequences, Biochemistry 35:7308-7315). When using poly(dIdC:dIdC) as the

substrate, GC-box b and GC-box b^{MET} showed noncompetitive and uncompetitive inhibition patterns, respectively (Figures 13 and 14). Both patterns require that the inhibitors and poly(dI-dC:dI-dC) bind to distinct sites on the enzyme surface. An uncompetitive pattern for GC-box b^{MET} suggests that potent inhibition is through an allosteric site and, in conjunction with the competitive inhibition with AdoMet, strongly implies inhibitor binding to the DCMTase:poly(dI-dC:dI-dC) complex in an ordered Bi-Bi mechanism (Spector & Cleland, 1981, Meanings of K_i for conventional and alternative-substrate inhibitors, Biochem. Pharm. 30:107). The noncompetitive inhibition pattern observed with GC-box b may result through weaker binding at the proposed allosteric site as well as binding at the active site. It is further speculated that the allosteric site is the same site where substrate inhibition originates. Because GC-box b and GC-box b^{MET} differ only in the methylation state of the single CpG, the 200-fold increased inhibition by GC-box b^{MET} is derived from the presence of the methyl group. Whether potent inhibition is caused by the methyl group itself or by greater DNA structural differences is not known.

Knowing that the DCMTase proceeds through the catalytic cycle in an ordered Bi-Bi mechanism allows for the determination of K_i , the dissociation constant for GC-box b^{MET} from the DCMTase:poly(dI-dC:dI-dC) complex (Spector & Cleland, 1981, Meanings of K_i for conventional and alternative-substrate inhibitors, Biochem. Pharm. 30:1-7). K_{ii} and K_{is} are conditional and can vary, thus K_i is the proper comparative. It is related to K_{ii} by this relation: $K_{ii} = K_i (1 + [\text{AdoMet}]/K_m^{\text{AdoMet}})$. Solving for K_i using the experimental data from Figure 14 it is found that $K_i = 2.5$ nM, a value about 10-fold lower than catalytic inhibition constant K_i .

Conclusion

Regulation of DNA replication and transcriptional activation by single-stranded DNA is known to occur (Takai, T., et al., 1994, Molecular cloning of MSSP-2, a c-myc gene single-strand binding protein: characterization of binding specificity and DNA replication activity, Nucleic Acids Res. 22:55776-5581; Rajavashisth, T.B., et al., 1989, Identification of a zinc finger protein that binds to the sterol regulatory element, Science 245:640-643; Tomonaga, T., & Levens, D., 1996, Activating transcription from single stranded DNA, Proc. Natl. Acad. Sci. USA 93:5830-5835). Nucleic acid regulation of DCMTase activity has previously been demonstrated. However, the requirement for micromolar concentrations of the polynucleic acids studied by Bolden

- et al. (1984, DNA methylation. Inhibition of *de novo* and maintenance methylation *in vitro* by RNA and synthetic polynucleotides, J. Biol. Chem. 259:12437-12443) to inhibit DCMTase implicates poor binding to the same site suggested in our studies with GC-box b^{MET}, or direct binding at the active site as competitive inhibitors. A stimulatory, *cis*-regulation by methylated CpG sites was reported to occur within single-stranded DNA using crude extracts (Christman, J.K., et al., 1995, 5-Methyl-2'-deoxycytidine in single-stranded DNA can act in *cis* to signal *de novo* DNA methylation, Proc. Natl. Acad. Sci. USA 92:7347-7351). While the mechanisms of regulation remain obscure in these cases, it is clear that they are distinct from the inhibition described herein. As previously stated, synthetic peptides mimicking portions of the DCMTase amino-terminus have been shown to gel mobility shift double-stranded DNA (Chuang, L.S., et al., 1996, Characterisation of independent DNA and multiple Zn-binding domains at the N terminus of human DNA-(cytosine-5) methyltransferase: modulating the property of a DNA-binding domain by contiguous Zn-binding motifs, Chia, J., and Li, B.F.L, J. Mol. Biol. 257:935-948). Although single-stranded DNA was apparently not studied, it would be interesting to systematically test these polypeptides for single-stranded DNA binding with and without methylated CpG dinucleotides.
- 20 The major finding of this Example concerns the identification of a second nucleic acid binding site that modulates the activity of DCMTase. Both the substrate inhibition studies and the dead-end inhibition studies with CD-box b^{MET} provide strong evidence for the existence of an allosteric site on the DCMTase surface. The kinetic studies demarcate the "allosteric" site, which is necessarily different from the "active" site where catalysis occurs. The novelty of these findings are drawn from the mechanistic insights that define the workings of the enzyme and the modulator in ways that have not been accessible to previous investigators.
- 30 GC-box b^{MET} is distinct in form and function from previously described DCMTase inhibitors. There is a need for DCMTase inhibitors that are not incorporated into DNA and that are mechanistically unlike 5-azadeoxycytidine (Belinsky, S.A., et al., 1996, Increased cytosine DNA-methyltransferase activity is target-cell-specific and an early event in lung cancer, Proc. Natl. Acad. Sci. USA 93:4045-4050; Szyf, M., 1996, The DNA methylation machinery as a target for anticancer therapy, Pharmacol. Ther. 70:1-37; Jones, P.A., 1996, DNA methylation errors and cancer, Cancer Res. 56:2463-2467). GC-box b^{MET} clearly interacts with a region of the enzyme that is

distinct from the active site and is highly sensitive to the presence of 5-methyl cytosine. The modulator described herein is a reversible antagonist of DCMTase function that provides a new class of therapeutics for treating developmental disorders such as cancer.

5

Example 3: Anti-proliferative Effects of DCMTase Inhibitors on Cells

It has been observed on several occasions that incubating mouse erythroleukemia (MEL) cells with GC-box b^{MET}, GC-box p and GC-box p^{MET} slows down the growth rate. The effect was shown to be concentration dependent. Inhibitor induced anti-proliferation was greatest at a concentration of 10 micromolar, 1 micromolar produced a moderate effect and 0.1 micromolar concentrations produced only a small difference in growth rate in comparison to untreated cells. As observed under the light microscope, concentrations of GC-box p^{MET} and GC-box p exceeding 2.5 micromolar induced MEL cells to produce small refractory particles of unknown content. GC-box b^{MET} also was observed to produce these particles at similar concentrations. The decrease in growth rate became more apparent after six days and three passages of the cells to fresh media containing the same inhibitor concentrations. Also, larger cells began to populate the culture after three days in a similar concentration-dependent manner. These large cells contained multiple nuclei and increased in number as length of incubation increased. After five days of incubation with 10 micromolar GC-box p^{MET}, the large multi-nucleate cells were observed to occur at about one in fifty regularly sized cells. Large multi-nucleate cells were also induced using the DCMTase anti-sense phosphorothioate used by Ramachandani et al., 1997 at a concentration of 10 micromolar.

-60-

SEQUENCE LISTING

(1) GENERAL INFORMATION

- (i) APPLICANT: Reich, Norbert O.
Flynn, James
- (ii) TITLE OF THE INVENTION: MODULATORS OF DNA CYTOSINE-5
METHYLTRANSFERASE AND METHODS FOR USE THEREOF
- (iii) NUMBER OF SEQUENCES: 110
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Merchant, Gould, Smith, Edell, Welter & Schmidt
 - (B) STREET: 11150 Santa Monica Boulevard, Suite 400
 - (C) CITY: Los Angeles
 - (D) STATE: CA
 - (E) COUNTRY: USA
 - (F) ZIP: 90025
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Diskette
 - (B) COMPUTER: IBM Compatible
 - (C) OPERATING SYSTEM: DOS
 - (D) SOFTWARE: FastSEQ for Windows Version 2.0
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE: 12-JUN-1998
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 60/057,411
 - (B) FILING DATE: 29-AUG-1997
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Canady, Karen S
 - (B) REGISTRATION NUMBER: 39,927
 - (C) REFERENCE/DOCKET NUMBER: 30794.30WO01
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 310 445-1140
 - (B) TELEFAX: 310 445-9031
 - (C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 26 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GGGGGGGRRK KGCGKGGKKGK KGKKGG

26

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 26 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

-61-

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

KGGRKKRDDD KRCGKRDDKK KKKKKG

26

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GGGGCGGGGC

10

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CCAACGTTGG

10

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CCAAGATTGG

10

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CGATCGATCG

10

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

-62-

CCAACGTTGG

10

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CCGTACGTAC GG

12

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GGGAATTCAA GGGGCGGGGC AAGGATCCAG

30

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CTGGATCCTT GCCCCGCCCC TTGAATCCC

30

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GGGAATTCAA ATGACGTCAA AAGGATCCAG

30

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

CTGGATCCTT TTGACGTCAT TTGAATCCC

30

(2) INFORMATION FOR SEQ ID NO:13:

-63-

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 50 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

CCTACCCACC CTGGATCCTT GCCCGCCCC TTGAATCCC AACCTCCAC

50

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

ATCCTTGCCC CGCCCCTTGA AT

22

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 14 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

TTGCCCCGCC CCTT

14

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 66 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GGGAATTCAT GGATCCTAAA NNNNNNNNNN NNGCNNNNNN NNNNNNTTTC AAGCTTGTGA
ATTCCC

60

66

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 66 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

CCCTTAAGTA CCTAGGATTT NNNNNNNNNN NNGCNNNNNN NNNNNNAAAG TTCGAACACT
TAAGGG

60

66

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

-64-

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GTGGGATGGG AACGAGTTGA GGAGGG

26

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

AGTGGTATGT ATCGATTATA GTTGGG

26

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GGAGGAAGTT TACGTATGGT ATGGGG

26

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

TGGGAGGGGA TTCGAGGTGA GAGTTG

26

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

ATAAAGTATT AGCGTAAGAG ATGAAG

26

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

-65-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

TGGAGGAGTT TACGGTGTA TGTGTT

26

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GGAGTAGGTA GACGTTAAGT ATGATG

26 GACGTTAAGT

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

GTGGGAAGGG GACGAATTTG AAGGTG

26 GACGAATTTG

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

TGTAATGTA TTCGTAAATG TAAGGG

26

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

TAATAGGGGA GACGTAAATG TAAGGG

26 GACGTAAATG

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

-66-

GAGTGTAGAA GTCGTAATAG ATTTAG

26

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

TGAGTAGGAA AGCGAAGAGG TGGTGG

26

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

TAGGTATTGG GCGGAAGGT GGGTGG

26

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

GGGGGTATAA TACGGTGTG GTAGGG

26

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

GGGTTGGGGT TTCGTGTGGG GGGTGT

26

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

TGTGGGTATG GCGGTGATA GTGAAG

26

(2) INFORMATION FOR SEQ ID NO:34:

-67-

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

GGATGATGGG GTCGAGAGTG GTGCTG

26

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

TAGTGGGTGG AGCGAGTGCT GGTGG

26

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

AGGGTGGGTG GCGGAGTTG TTGTTG

26

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

GTGAGGAGGG AGCGGGAATG GGGGTG

26

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

GGGGGTGGGG AGCGGAGGGG GGTGAG

26

(2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double

-68-

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

TGTTGGAGGG GGCGAAGGTG GTTTTG

26

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

GGGGGGGGGG GGCGAGGGGT AGATGG

26

(2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

GGGGGAGGGG TTCGGTGATA GGTAGG

26

(2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

GGGGGGGGGG TACGTGGGAT GGTATG

26

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

GTGTAGGGAG TGCGAGGGGG TGTAAG

26

(2) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

-69-

GGGGGGGGGT AGCGGTTAGA TGGTGG

26

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

GGGGTGAGGG GCGGGGGTT AGTGGG

26

(2) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

GAGGGGGGGT TCGTAGGGG GGTGGG

26

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

TGTGGAGGTG GCGGGGAAAG GTGATG

26

(2) INFORMATION FOR SEQ ID NO:48:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

GGGGGGATGG GACGGATGGG GGGGGG

26

(2) INFORMATION FOR SEQ ID NO:49:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

GGGGGTGGGG TGCGAGAGAG TTGGGG

26

(2) INFORMATION FOR SEQ ID NO:50:

-70-

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

GAGGGGTGGA GCGGAGGTG GGTGG

26

(2) INFORMATION FOR SEQ ID NO:51:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

GGGGGGGGG GCGGATAAG GGTGTG

26

(2) INFORMATION FOR SEQ ID NO:52:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

TGGGGGGGGG GCGGGGGGA GTTTGA

26

(2) INFORMATION FOR SEQ ID NO:53:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

GGGGGGGAGG GCGGATAGT TGTGTG

26

(2) INFORMATION FOR SEQ ID NO:54:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

GGGTGGGGTG GCGGTGGGG TGTGGG

26

(2) INFORMATION FOR SEQ ID NO:55:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid

-71-

- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

GAGGGGGGGG AGCGGAGGGG GTTGGG

26

(2) INFORMATION FOR SEQ ID NO:56:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 26 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

GGGGGGGAAG GCGGTGGGGT TGGGTG

26

(2) INFORMATION FOR SEQ ID NO:57:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

GGGAGGGGGG GCGATGGGGT GGTGG

25

(2) INFORMATION FOR SEQ ID NO:58:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 26 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

GGGTGGGGGT GCGGTTGTGG GTGGGG

26

(2) INFORMATION FOR SEQ ID NO:59:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 26 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

GGGAGGGGGT GCGGTTGGGT ATGTGG

26

(2) INFORMATION FOR SEQ ID NO:60:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 26 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

-72-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

GGGGAGGGTG GCGGGGTATG GAGTGG

26

(2) INFORMATION FOR SEQ ID NO:61:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

GGGGGGGGAG TGCCTTGATG GGTGTG

26

(2) INFORMATION FOR SEQ ID NO:62:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

GGGGGGGTGG ATCGTGGGGG GAGGGG

26

(2) INFORMATION FOR SEQ ID NO:63:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

GGGGTAGGGT GCGGGGGGGG GTATGG

26

(2) INFORMATION FOR SEQ ID NO:64:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

GGGATGGGGG TGCGGGGTAT GGGGGG

26

(2) INFORMATION FOR SEQ ID NO:65:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

GGGAGGGGGT AGCGGGAGTG TGTGTG

26

-73-

(2) INFORMATION FOR SEQ ID NO:66:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

GGGGGTAAGG GGCCTAAGAA TGGGGG

26

(2) INFORMATION FOR SEQ ID NO:67:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

GGGGGGGTGG TTCGTAAATG GGGGGT

26

(2) INFORMATION FOR SEQ ID NO:68:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

GGTGGGAGAG GCGGTGGTGT AGGTAG

26

(2) INFORMATION FOR SEQ ID NO:69:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

GGGGGGGGTG TACGAGGTTT GTGTGG

26

(2) INFORMATION FOR SEQ ID NO:70:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

TGGTGGAGGG GGCGAAGAAG TGTGTG

26

(2) INFORMATION FOR SEQ ID NO:71:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs

-74-

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

GGGGGTGGGA TGCGGAATAA GGATGG

26

(2) INFORMATION FOR SEQ ID NO:72:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

TGAGGGGGAG GGCGAATAGA TGGTGG

26

(2) INFORMATION FOR SEQ ID NO:73:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

GGGGGGAGTA AGCGGGGGTG TGGTGG

26

(2) INFORMATION FOR SEQ ID NO:74:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

TGAAGGGGGG TGCGGGGTGT GGGGG

25

(2) INFORMATION FOR SEQ ID NO:75:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:

GTGGTGATGG GGCGGGGTGG TAGTGG

26

(2) INFORMATION FOR SEQ ID NO:76:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

-75-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

TGGAGGGGTA GCGGTGGGGT GATGGG

26

(2) INFORMATION FOR SEQ ID NO:77:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

GGTAGGGAGT GCGGGTGGT GATGGG

26

(2) INFORMATION FOR SEQ ID NO:78:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:

GGGTGTAGAG GCGGGGAGTA GAGGGG

26

(2) INFORMATION FOR SEQ ID NO:79:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:

GGGTGGGTTT GCGTAATTG TGTGGG

26

(2) INFORMATION FOR SEQ ID NO:80:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:

GGGTGTGTTG GCGGTGGGGT ATGTAG

26

(2) INFORMATION FOR SEQ ID NO:81:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:

TGGGGAGAAT GCGGGGGGGT GGTGGG

26

-76-

(2) INFORMATION FOR SEQ ID NO:82:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:82:

TATGGTGGGA GCGGGGGGG GGTGG

26

(2) INFORMATION FOR SEQ ID NO:83:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:83:

TGGGGAAAGA GCGTGAGTG GGGGGG

26

(2) INFORMATION FOR SEQ ID NO:84:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:84:

TGTAGGGGAG GACGGGGGAT GGGGTG

26

(2) INFORMATION FOR SEQ ID NO:85:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:85:

GGGTGGGTAA TCGTAGGGT GGGGGG

26

(2) INFORMATION FOR SEQ ID NO:86:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:86:

GTGTGGGTAA GCGGTATGG GGTGG

26

(2) INFORMATION FOR SEQ ID NO:87:

(i) SEQUENCE CHARACTERISTICS:

-77-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:87:

TGGAGGGTGT TCGGGTGAGG TGGTGG

26

(2) INFORMATION FOR SEQ ID NO:88:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:88:

GGTGGTGGTG ATCGGGGTTG TGATGG

26

(2) INFORMATION FOR SEQ ID NO:89:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:89:

GGGGGTAAAG TCGGGGTGGT TGATGG

26

(2) INFORMATION FOR SEQ ID NO:90:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:90:

GTGGAGGTGT TCGGTAGTGT GGGAGG

26

(2) INFORMATION FOR SEQ ID NO:91:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:91:

GTGGGGAATG GTCGGTTATG GTGGGG

26

(2) INFORMATION FOR SEQ ID NO:92:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:92:

-78-

GGGATGTGGT AGCGGGGGTG TGTTAG

26

(2) INFORMATION FOR SEQ ID NO:93:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:93:

GGGGTAGGAG TTCGTAGGGG TGTGTT

26

(2) INFORMATION FOR SEQ ID NO:94:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:94:

GAGGTGGTGG ATCGGGATGA TGGATT

26

(2) INFORMATION FOR SEQ ID NO:95:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:95:

TGGGGGGAAA TACGGGGAGG GTGGTA

26

(2) INFORMATION FOR SEQ ID NO:96:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:96:

GGAGTAGGGT TACGTGGTGG TAATGG

26

(2) INFORMATION FOR SEQ ID NO:97:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:97:

GAGGAGTAAA GGCGTGTGTT GTGGTG

26

(2) INFORMATION FOR SEQ ID NO:98:

-79-

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:98:

TGGATGAGAG TGGGTGTATG ATAAGG

26

(2) INFORMATION FOR SEQ ID NO:99:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:99:

AGGGTTAGTG AACGGGGGGG AGGTGG

26

(2) INFORMATION FOR SEQ ID NO:100:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:100:

GAGAAGGGTA AACGTGGGGG AGGGGA

26

(2) INFORMATION FOR SEQ ID NO:101:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:101:

TGGGGGGGGG GCGGGGGGA GTTTGA

26

(2) INFORMATION FOR SEQ ID NO:102:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 56 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:102:

GAACAATGGG GCGCTGGGG GGGGGGCGG GGGGCTTTA GCTATGTCAG AATTCA

56

(2) INFORMATION FOR SEQ ID NO:103:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double

-80-

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:103:

GGGATGGGGG TGCGGGGTAT GGGGGG

26

(2) INFORMATION FOR SEQ ID NO:104:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 57 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:104:

GGGGAACAGC GAGCACCGAA GGGGGTGCGG GGTATGGGAG GGTCCCCGGG CTTGAGC

57

(2) INFORMATION FOR SEQ ID NO:105:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:105:

GGTGGTGGTG ATCGGGGTTG TGATGG

26

(2) INFORMATION FOR SEQ ID NO:106:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 56 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:106:

TGTCCTTCTT GTGGTGGTGG TAGAGGTCGT GGTGTGATG GTGGCTCGGT GTGTGT

56

(2) INFORMATION FOR SEQ ID NO:107:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:107:

GAGGGGGGGG AGCGGAGGGG GTTGGG

26

(2) INFORMATION FOR SEQ ID NO:108:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 56 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:108:

-81-

GAGGGGTGTA GCCGCGAGGG GGCGGAGCGG AGGGGGAGGG CCCTGGTCCC GCCGCC

56

(2) INFORMATION FOR SEQ ID NO:109:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:109:

CCCCACCCAC AACGCCACCC CCACCC

26

(2) INFORMATION FOR SEQ ID NO:110:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 56 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:110:

TCTTTAAATG GTGCGGTCCA CCCCCACCGC CACCCCCACC CCCCACTGGA GCAAGG

56

What is claimed is:

1. A synthetic oligonucleotide comprising a C-5 methylcytosine and which recognizes and binds an allosteric site on DNA cytosine methyltransferase (DCMTase) thereby modulating DCMTase activity associated with the allosteric site.
5
2. The synthetic oligonucleotide of claim 1, wherein the modulating comprises inhibition.
3. The synthetic oligonucleotide of claim 1, wherein the modulating comprises activation.
10
4. The synthetic oligonucleotide of claim 1, wherein the C-5 methylcytosine is present as a 5mCpG dinucleotide.
5. The synthetic oligonucleotide of claim 1, wherein the DCMTase is from a mammal, bird, fish, amphibian, reptile, insect, plant or fungus.
- 15 6. The synthetic oligonucleotide of claim 5, wherein the mammal is selected from the group consisting of mouse and human.
7. The synthetic oligonucleotide of claim 1 having an inhibition constant of not greater than 1000 nM.
8. The synthetic oligonucleotide of claim 7 having an inhibition constant of not greater than 200 nM.
20
9. The synthetic oligonucleotide of claim 8 having an inhibition constant of not greater than 20 nM.
10. The synthetic oligonucleotide of claim 1 comprising a nucleotide sequence as shown in Figure 1B and designated GC-box b^{MET} (SEQ ID NO:10), GC-box p^{MET} (SEQ ID NO:10), GC-box c^{MET} (SEQ ID NO:13), GC-box d^{MET} (SEQ ID NO:14), GC-box e^{MET} (SEQ ID NO:15), or CRE a^{MET} (SEQ ID NO:11).
25
11. A method of inhibiting methylation of DNA comprising contacting a DCMTase with a synthetic inhibitor molecule so as to form an enzyme/synthetic inhibitor molecule complex in the presence of the DNA, wherein the synthetic inhibitor

molecule comprises a C-5 methylcytosine which recognizes and binds an allosteric site on DCMTase, thereby inhibiting DNA methyltransferase activity.

12. A method of inhibiting proliferation of cancer cells comprising administering to a subject a synthetic inhibitor molecule which recognizes and binds an allosteric site on DCMTase thereby resulting in an enzyme/synthetic inhibitor molecule complex, the presence of the complex inhibiting DCMTase-mediated methylation of DNA, thereby inhibiting proliferation of the cancer cells.
13. The method of claim 12, wherein the cancer cell is from lung, breast, prostate, pancreas or colon.
14. The method of claim 11 or 12, wherein the synthetic inhibitor molecule is an oligonucleotide of any one of claims 1-10.
15. The method of claim 12, 13, or 14, wherein the subject is a human.
16. The method of claim 12, 13, or 14, wherein the subject is an animal.
17. The method of claim 16, wherein the animal is porcine, piscine, avian, feline, equine, bovine, ovine, caprine or canine.
18. A method of identifying a molecule which recognizes and binds an allosteric site on DCMTase comprising:
 - (a) contacting a molecule with DCMTase in the presence of DNA and AdoMet;
 - (b) measuring DCMTase activity, an increase or decrease in DCMTase activity being indicative of a modulator of DCMTase; and
 - (c) determining whether the modulation of DCMTase activity is via binding an allosteric site on DCMTase.
19. The method of claim 18, wherein the modulator is an inhibitor.
20. The method of claim 18, wherein DCMTase activity is measured using a steady-state assay.

FIG. 1a.

Synthetic DNA Substrates Mimicking Transcriptional
Cis-Regulatory Elements


GC-box a: 5'-GGGAATTCAAGGGGCGGCAAGGATCCAG -3'
GC-box b: 5'-CTGGATCCTTGCCCGGCCCTTGAATTCCC -3'
GC-box b MET: 5'-CTGGATCCTTGCCC^mCGCCCCCTTGAATTCCC -3'
CRE a: 5'-GGGAATTCAAATGACGTCAAAGGATCCAG -3'
CRE b: 5'-CTGGATCCTTTTGACGTCATTGAATTCCC -3'
CRE a MET: 5'-GGGAATTCAAATGA^mCGTCAAAGGATCCAG -3'

FIG. 1b.

NAME	NUCLEO- TIDES	Sequence	Kil IC50 (nM) (nM)
GC-Box b (SEQ ID NO: 10)	30	5'-CTGGATCCTTGCCCCGCCCCCTTGAATTCCC-3'	6800
GC-Box bMET (SEQ ID NO: 10)	30	5'-CTGGATCCTTGCCCCmCGCCCCCTTGAATTCCC-3'	20 15
GC Box pMET (SEQ ID NO: 10)	30	5'-CTGGATCCTTGCCCCmCGCCCCCTTGAATTCCC-3'	5
GC-Box cMET (SEQ ID NO: 13)	50	5'- CCTACCCACCCTGGATCCTTGCCCCmCGCCCCCTTGAATTCCCCAACCCCTCCAC-3'	30
GC Box dMET (SEQ ID NO: 14)	22	5'-ATCCTTGCCCCmCGCCCCCTTGAAT-3'	50
GC-Box eMET (SEQ ID NO: 15)	14	5'TTGCCCCmCGCCCCCTT-3'	150
CRE aMET (SEQ ID NO: 11)	30	5'-GGGAATTCAAATGAmCGTCAAAAGGATCCAG-3'	> 300


3/26

FIG. 2.




4/26

FIG. 3.




5/26

FIG. 4



6/26

FIG. 5



7/26

FIG. 6.

Primer C

5'-GGGAATTCATGGATCCTAAANNNNNNNNNNCGNNNNNNNNNTTCAAGCTTGTGAATTCCC-3'
3'-CCCTTAAGTACCTAGGATTTNNNNNNNNNNGCGNNNNNNNNNAAAGTTTCGAACACTTAAGGG-5'

Primer D

FIG.7a.

8/26

STARTING POPULATION

GTGGGATGGGAACGAGTTGAGGAGGG
AGTGGTATGTATCGATTATACGTTGGG
GGAGGAAGTTTACGTATGGTATGGGG
TGGGAGGGGATTGAGGTGAGAGTTG
ATAAAGTATTAGCGTAAGAGATGAAG
TGGAGGAGTTTACGGTGTAATTGTTT
GGAGTAGGTAGACGTTAAGTATGATG
GTGGGAAGGGGACGAATTTGAAGGTG
TGGTAATGTATTGTAATGTAAGGG
TAATAGGGGAGACGTAAATGTAAGGG
GAGTGTAGAAGTCGTAATAGATTTAG
TGAGTAGGAAAGCGAAGAGGTGTTGG

FIG.7b.

GENERATION 1

TAGGTATTGGGGCGGAAGGTGGGTGG
GGGGGTATAATACGGTGTTGGTAGGG
GGGTTGGGGTTTCGTGTGGGGGGTGT
TGTGGGTATGGGCGGTGATAGTGAAG
GGATGATGGGGTCGAGAGTGGTGGTG
TAGTGGGTGGAGCGAGTGGTGGTTGG
AGGGTGGGTGGGCGGAGTTGTTGTTG
GTGAGGAGGGAGCGGGAATGGGGGTG
GGGGGTGGGGAGCGGAGGGGGGTGAG
TGTTGGAGGGGGCGAAGGTGGTTTTG

FIG.7c.

GENERATION 3

GGGGGGGGGGGGCGAGGGGTAGATGG
GGGGGAGGGGTTGCGGTGATAGGTAGG
GGGGGGGGGGGTACGTGGGATGGTATG
GTGTAGGGAGTGCGAGGGGGGTGTAAG
GGGGGGGGGGTAGCGGTTAGATGGTGG
GGGGTGAGGGGGCGGGGGTTAGTGGG
GAGGGGGGGGTTGCGTAGGGGGGTGGG
TGTGGAGGTGGGCGGGAAAGGTGATG
GGGGGGATGGGACGGATGGGGGGGGG
GGGGGTGGGGTGCGAGAGAGTTGGGG
GAGGGGTGGAGGCGGAGGTGGGTTGG
GGGGGGGGGGGGCGATAAGGGTGTG

9/26

FIG.7d.

G#	GpT	TpG	GENERATION 5	TpG	GpT	G#
11	.	.	TGGGGGGGGG <u>CGGGG</u> AGTTGA	.	.	7
11			GGGGGGGAGGGG <u>CGG</u> ATAGTTGTGTG	5
10	GGGTGGGTGGG <u>CGG</u> TGGGTGTGGG	9
10			GAGGGGGGGA <u>GCGG</u> AGGGGTTGGG	.	.	9
10			GGGGGGGAAGGG <u>CGTGGG</u> TTGGGTG	8
10			-GGAGGGGGG <u>CGG</u> ATGGGTGGTGG	8
10	GGGTGGGGTGG <u>CGT</u> TGTGGGTGGG	8
10	.	.	GGAGGGGTGG <u>CGG</u> TGGGTATGTGG	7
10	.	.	GGGAGGGTGGG <u>CGG</u> ATGGAGTGG	7
10	.	.	GGGGGGGAGT <u>GCGT</u> GATGGGTGTG	6
9	.	.	GGGGGGGTGGAT <u>CGTGGG</u> GAGGGG	.		10
9	GGGTAGGTGGG <u>CGG</u> GGGGGTATGG	.	.	9
9	.	..	GGGATGGGGT <u>GCGG</u> GATGGGGG	.	.	9

10/26

FIG. 7e.

G#	GpT	TpG	GENERATION 5	TpG	GpT	G#
9	.	.	GGGAGGGGTAGCGG <u>G</u> AGTGTGTG	7
9	.	.	GGGGCTAAGGGCGT <u>A</u> GAATGGGG	.		6
9	..	.	GGGGGGTGGTTCGGT <u>A</u> ATGGGGGT	.	..	7
9	.	.	GGTGGAGAGGGCGTGGTGTAGGTAG	6
9	..	.	GGGGGGGTGTACGAGGTTTGTGTGG	6
9	.	..	TGGTGGAGGGGCGAAGAAAGTGTGTG	5
9	.	..	GGGGGTGGGATGCGGAATAAGGATGG	.		6
9	.	.	TGAGGGGAGGGCGAATAGATGGTGG	..	.	7
8	.	.	GGGGGGAGTAAGCGGGGTGTGTGG	9
8	.	..	TGAAGGGGGTGC <u>G</u> GGGTGTGGGGG-	9
8	GTGGTGTGGGGCGGGGTGGTAGTGG	8
8	.	.	TGGAGGGGTAGGCGTGGGGTGTGGG	8

11/26

FIG. 7f.

GENERATION 5						
G#	GpT	TpG	TpG	GpT	G#	
8	8	GGTAGGAGTG <u>GC</u> GGTGGTGA <u>TGGG</u>
8	8	GGGTGTAGAGGCG <u>CGG</u> AGTAGAGGGG
8	7	GGGTGGGTTTGGC <u>CGT</u> AATTGTGTGGG
8	6	GGGTGTGTGGC <u>CGT</u> GGGGTATGTAG
7	10	TGGGGAGAAATGCG <u>CGG</u> GGGTGGTGGG
7	10	TATGTTGGGAGGCG <u>CGG</u> GGGGGGGTTGG
7	9	TGGGGAAAGAGGCG <u>CGT</u> AGTGGGGGGG
7	9	TGTAGGGAGGACG <u>GGG</u> GATGGGGTG
7	9	GGGTGGGTAATGCGT <u>AGG</u> GTGGGGGG
7	8	GTGTGGGTAAGGCGGTA <u>TGG</u> GGGTGG
7	8	TGGAGGGTGTGCGGTA <u>GAG</u> GTGGTGG
7	7	GGTGGTGTGATCGGCGT <u>TGT</u> GATGG

12/26

FIG. 7g.

GENERATION 5				
G#	GpT	TpG	GpT	G#
7	7
7	7
7	7
7	7
7	6
7	5
6	8
6	6
6	6
6	4
5	10
5	9

13/26

FIG.8a.




FIG.8b.

	-12	-11	-10	-9	-8	-7	-6	-5	-4	-3	-2	-1	C	G	1	2	3	4	5	6	7	8	9	10	11	12
%G	73	82	84	61	71	73	73	59	49	57	53	76	0	100	57	61	65	53	61	55	51	63	59	53	73	88
%A	2	10	6	22	10	10	22	20	29	20	18	12	0	0	10	20	18	20	10	12	12	14	12	6	6	6
%T	25	8	10	16	18	16	4	20	22	22	29	12	0	0	33	18	16	27	29	33	37	22	29	41	20	6
%C	0	0	0	0	0	0	0	0	0	0	0	0	100	0	0	0	0	0	0	0	0	0	0	0	0	0

14/26

FIG.9a

DEFINITION Lyt-2.2 gene, T- cell differentiation antigen, 3' UTR.
 ACCESSION GB_RO:MMLYT22

GAACAATGGGGCGCTGGGGGGGGGGCGGGGGGAGTTTGA 5100 5110 5120 5130 5140
 TGGGGGGGGGGCGGGGGGAGTTTGA
 |||||
 |||||

DEFINITION homeo box 2.6 (Hox-2.6) mRNA
 ACCESSION GB_RO:MUSHOX26

GGGATGGGGGTGCGGGGTATGGGGG
 |||||
 GGGGAACAGCGAGCACCGAAGGGGTGCGGGGTATGGGAGGGTCCCGGGCTTGAGC 870 880 890 900 910 920

DEFINITION growth arrest-specific promoter gene, gas-1
 ACCESSION GB_RO:MMGAS1PRA

GGTGGTGGTGATCGGGGTGTGATGG
 |||||
 TGTCCTTCTTGTGGTGGTAGAGGTCGTGTTGTGATGGTGCTCGGTGTGTGT 2480 2490 2500 2510 2520

15/26

FIG. 9b.

DEFINITION pim-1 proto-oncogene, pim-1 protein kinase, CpG island,
5' UTR region.

ACCESSION GB_RO:MUSPIM1

GAGGGGGGAGCGGAGGGGTTGGG
| | | | | | | | | | | | | | | | | |
GAGGGGTGTAGCCGCGAGGGGCGGAGCGGAGGGGAGGGCCCTGTCCCGCCGCC
1500 1510 1520 1530 1540

DEFINITION neuronal dihydropyridine-sensitive L-type calcium
channel alpha-1 subunit mRNA, 3' UTR.

ACCESSION GB_RO:MUSDHPCC

CCCCACCCACAACGCCACCCCCACCC
| | | | | | | | | | | | | | | | | |
TCCTTAAATGGTGCGGTCCACCCCCACCGCCACCCCCCACTGGAGCAAGG
8330 8340 8350 8360 8370 8380

16/26

FIG. 9c.

HUMAN SEQUENCES

DEFINITION Huntington's Disease Region, chromosome 4p16.3.
ACCESSION GB_PR:HSL1C2

DEFINITION Human Down Syndrome region of chromosome 21.
ACCESSION GB_HTG:HSAC000002

DEFINITION upstream region of HoxA7 gene, CpG island.
ACCESSION GB_PR:HSHCRDNA

DEFINITION chromosome 22 CpG island DNA
ACCESSION GB_PR:HS303B3

DEFINITION CpG island DNA.
ACCESSION GB_PR:HS167B9F

DEFINITION Y chromosome sex determining region, Y p pseudoautosomal
 boundary, PAB1.
ACCESSION GB_PR:HSCAMF3X1

DEFINITION creatine transporter and paralogous genes, pericentomeric
 repeats on chromosome 16.
ACCESSION GB_PR:HSU41302

DEFINITION cathepsin D (cat D) gene, exon 5.
ACCESSION GB_PR:HUMCATD3

17/26

FIG.9d.

DEFINITION	argininosuccinate synthetase gene 5' end, CpG island
ACCESSION	GB_PR:HSASG5E
DEFINITION	argininosuccinate synthetase gene 5' end, CpG island
ACCESSION	GB_PR:HUMAS1
DEFINITION	vimentin gene, 5' regulatory region, CpG island.
ACCESSION	GB_PR:HUMVIM
DEFINITION	vimentin gene, exon 1, 5' end CpG island.
ACCESSION	GB_PR:HUMVIM02
DEFINITION	vimentin gene, 5' end, CpG island.
ACCESSION	GB_PR:HUMVIMAA
DEFINITION	vimentin gene, 5' end, CpG island
ACCESSION	GB_PR:HSVIM5RR
DEFINITION	chromosome 22 DNA *SEQUENCING IN PROCESS*, CpG island
ACCESSION	GB_HTG:HS170A21

18/26

FIG. 10.






FIG. 13.



19/26



20/26

FIG.12a.





FIG.12b.



21/26

FIG.14.






FIG.16.



22/26

FIG. 15.



23/26

FIG. 17a.




FIG. 17b.





FIG. 17c.



24/26

FIG. 18.




FIG. 19a.





FIG. 19b.



SUBSTITUTE SHEET (RULE 26)

25/26

FIG.20.




FIG.21.




FIG.22.



26/26

FIG.23a.




FIG.23b.





FIG.23c.



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/12351

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :G01N 31/00

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : Please See Extra Sheet.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,578,716 A (SZYF et al) 26 November 1996, column 2, line 43 through column 8.	1-13,18-20



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

03 AUGUST 1998

Date of mailing of the international search report

24 SEP 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JAMES O. WILSON

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/12351

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☒ Claims Nos.: 14-17
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/12351

A. CLASSIFICATION OF SUBJECT MATTER:

US CL : 435/6; 514/43; 536/22.1, 23.1, 23.2, 25.32

B. FIELDS SEARCHED

Minimum documentation searched

Classification System: U.S.

435/6; 514/43; 536/22.1, 23.1, 23.2, 25.32

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

USPATFULL

WPIDS

CAS ONLINE

BIOSIS